


Energy Changes

Q1.

The figure below shows magnesium burning in air.



© Charles D Winters/Science Photo Library

- (a) Look at the figure above.

How can you tell that a chemical reaction is taking place?

(1)

- (b) Name the product from the reaction of magnesium in the figure.

(1)

- (c) The magnesium needed heating before it would react.

What conclusion can you draw from this?

Tick **one** box.

The reaction is reversible

The reaction has a high activation energy

The reaction is exothermic

Magnesium has a high melting point

(1)

- (d) A sample of the product from the reaction in the figure above was added to water and shaken.

Universal indicator was added.

The universal indicator turned blue.

What is the pH value of the solution?

Tick **one** box.

1

4

7

9

(1)

- (e) Why are nanoparticles effective in very small quantities?

Tick **one** box.

They are elements

They are highly reactive

They have a low melting point

They have a high surface area to volume ratio

(1)

- (f) Give **one** advantage of using nanoparticles in sun creams.

(1)

- (g) Give **one** disadvantage of using nanoparticles in sun creams.

(1)


- (h) A coarse particle has a diameter of 1×10^{-6} m.
A nanoparticle has a diameter of 1.6×10^{-9} m.

Calculate how many times bigger the diameter of the coarse particle is than the diameter of the nanoparticle.

(2)
(Total 9 marks)

Q2.

A student investigated simple cells using the apparatus shown in the figure below.




- If metal **2** is more reactive than metal **1** then the voltage measured is positive.
- If metal **1** is more reactive than metal **2** then the voltage measured is negative.
- The bigger the difference in reactivity of the two metals, the larger the voltage produced.

The student's results are shown in the table below.

Metal 1 \ Metal 2	Chromium	Copper	Iron	Tin	Zinc
Chromium	0.0 V				
Copper	1.2 V	0.0 V			
Iron	0.5 V	not measured	0.0 V		
Tin	0.8 V	-0.4 V	0.3 V	0.0 V	
Zinc	0.2 V	-1.0 V	-0.3 V	-0.6 V	0.0 V

- (a) The ionic equation for the reaction occurring at the zinc electrode in the simple cell made using copper and zinc electrodes is:



Zinc is oxidised in this reaction.

Give a reason why this is oxidation.

(1)

- (b) Look at the table above.

Which **one** of the metals used was the least reactive?

Give a reason for your answer.

Metal _____

Reason _____

(2)

- (c) Predict the voltage that would be obtained for a simple cell that has iron as metal **1** and copper as metal **2**.

Explain your answer.

(3)

(d) Hydrogen fuel cells have been developed for cars.

Write a word equation for the overall reaction that takes place in a hydrogen fuel cell.

(1)

(e) Write the **two** half equations for the reactions that occur at the electrodes in a hydrogen fuel cell.


(2)

(Total 9 marks)

Q3.

This question is about the reaction of ethene and bromine.

The equation for the reaction is:




(a) Complete the reaction profile in **Figure 1**.

Draw labelled arrows to show:

- The energy given out (ΔH)
- The activation energy.

Figure 1



(3)


- (b) When ethene reacts with bromine, energy is required to break covalent bonds in the molecules.

Explain how a covalent bond holds two atoms together.

(2)

- (c) **Figure 2** shows the displayed formulae for the reaction of ethene with bromine.

Figure 2




The bond enthalpies and the overall energy change are shown in the table below.

	C=C	C-H	C-C	C-Br	Overall energy change
Energy in kJ / mole	612	412	348	276	-95

Use the information in the table above and **Figure 2** to calculate the bond energy for the Br-Br bond.

- (d) **Figure 3** shows the reaction between ethene and chlorine and is similar to the reaction between ethene and bromine.

Figure 3



“The more energy levels (shells) of electrons an atom has, the weaker the covalent bonds that it forms.”

Use the above statement to predict and explain how the overall energy change for the reaction of ethene with chlorine will differ from the overall energy change for the reaction of ethene with bromine.

(6)


(Total 14 marks)

Q4.

This question is about hydrocarbons.

- (a) Most of the hydrocarbons in crude oil are alkanes.
- (i) Large alkane molecules can be cracked to produce more useful molecules.

The equation shows the cracking of dodecane.



Give **two** conditions used to crack large alkane molecules.

1. _____

2. _____

(2)

- (ii) The products hexene and ethene are alkenes.


Complete the sentence.

When alkenes react with bromine water the colour changes
from orange to _____ .

(1)

- (iii) Butane (C_4H_{10}) is an alkane.


Complete the displayed structure of butane.



(1)

- (b) A group of students investigated the energy released by the combustion of four hydrocarbon fuels.

The diagram below shows the apparatus used.



Each hydrocarbon fuel was burned for two minutes.

Table 1 shows the students' results.

Table 1

Name and formula of hydrocarbon fuel	After two minutes			Energy released by 1.0 g of fuel in kJ	Relative amount of smoke in the flame
	Mass of fuel used in g	Temperature increase of water in °C	Energy released by fuel in kJ		
Hexane, C_6H_{14}	0.81	40	16.80	20.74	very little smoke
Octane, C_8H_{18}	1.10	54	22.68	20.62	some smoke

Decane, C ₁₀ H ₂₂	1.20	58	24.36		smoky
Dodecane, C ₁₂ H ₂₆	1.41	67	28.14	19.96	very smoky

- (i) Calculate the energy released by 1.0 g of decane in kJ.

Energy released = _____ kJ

(2)

- (ii) Suggest **one** improvement to the apparatus, or the use of the apparatus, that would make the temperature increase of the water for each fuel more accurate.

Give a reason why this is an improvement.

(2)

- (iii) The students noticed that the bottom of the beaker became covered in a black substance when burning these fuels.

Name this black substance.

Suggest why it is produced.

(2)

- (iv) A student concluded that hexane is the best of the four fuels.

Give **two** reasons why the results in **Table 2** support this conclusion.

1. _____

2. _____

(2)

- (c) **In this question you will be assessed on using good English, organising information clearly and using specialist terms where appropriate.**

Most car engines use petrol as a fuel.

- Petrol is produced from the fractional distillation of crude oil.
- Crude oil is a mixture of hydrocarbons.
- Sulfur is an impurity in crude oil.

Car engines could be developed to burn hydrogen as a fuel.

- Hydrogen is produced from natural gas.
- Natural gas is mainly methane.

Table 2 shows information about petrol and hydrogen.

	Petrol	Hydrogen
State of fuel at room temperature	Liquid	Gas
Word equation for combustion of the fuel	petrol + oxygen \rightarrow carbon dioxide + water	hydrogen + oxygen \rightarrow water
Energy released from combustion of 1 g of the fuel	47 kJ	142 kJ

Describe the **advantages** and **disadvantages** of using hydrogen instead of petrol in car engines.

Use the information given and your knowledge and understanding to answer this question.


(6)
(Total 18 marks)

Q5.

This question is about temperature changes.

- (a) A student investigated the temperature change when 8 g of sodium nitrate dissolves in 50 cm³ of water.

The diagram below shows the apparatus the student used.



The student did the experiment five times.

Table 1 shows the results.

Table 1

Experiment	Decrease in temperature of water in °C
1	5.9
2	5.7
3	7.2
4	5.6
5	5.8

- (i) Calculate the mean decrease in temperature.
Do not use the anomalous result in your calculation.

Mean decrease in temperature = _____ °C

(2)

- (ii) Suggest **one** change in the apparatus in the diagram above which would improve the accuracy of the results.
Give a reason for your answer.

(2)

- (b) The student investigated the temperature change when different masses of sodium carbonate were added to 50 cm³ of water at 20 °C.

Table 2 below shows the results.

Table 2

Mass of sodium carbonate in g	Final temperature of solution in °C
2.0	21.5
4.0	23.0
6.0	24.5
8.0	26.0
10.0	26.6

12.0	26.6
14.0	26.6

Describe the relationship between the mass of sodium carbonate added and the final temperature of the solution.

Use values from **Table 2** in your answer.

(3)


(Total 7 marks)

Q6.

This question is about ammonia and fertilisers.

- (a) Ammonia is produced by a reversible reaction.

The equation for the reaction is:



Complete the sentence.

The forward reaction is exothermic, so the reverse reaction

is _____

(1)

- (b) Calculate the percentage by mass of nitrogen in ammonia (NH₃).

Relative atomic masses (A_r): H = 1; N = 14

You **must** show how you work out your answer.

Percentage by mass of nitrogen = _____ %

(3)

(c) A neutral solution can be produced when ammonia reacts with an acid.





(i) Give the pH of a neutral solution.

pH _____

(1)

(ii) Which of these ionic equations shows a neutralisation reaction?

Tick (✓) **one** box.



(1)

(iii) Name the salt produced when ammonia reacts with hydrochloric acid.

(1)




(d) **In this question you will be assessed on using good English, organising information clearly and using specialist terms where appropriate.**

Farmers use ammonium nitrate as a fertiliser for crops.

Rainwater dissolves ammonium nitrate in the soil.

Some of the dissolved ammonium nitrate runs off into rivers and lakes.

The graphs **A**, **B** and **C** below show information about the use of ammonium nitrate as a fertiliser. A hectare is a measurement of an area of land.



Suggest how much ammonium nitrate farmers should use per hectare.

Give reasons for your answer.

Use information from graphs **A**, **B** and **C**.

(6)
(Total 13 marks)

Q7.

This question is about energy changes in chemical reactions.


- (a) Complete the word equation for the combustion of hydrogen.



(1)

- (b) **Figure 1** shows a simple energy level diagram.

Figure 1



- (i) Which arrow, **A**, **B** or **C**, shows the activation energy?

Tick (✓) **one** box.

A

B



C



(1)

- (ii) What type of reaction is shown by the energy level diagram in **Figure 1**?
Give a reason for your answer.

Type of reaction _____

Reason _____

(2)

- (iii) For a reaction, the value of **A** is 1370 kJ and **C** is 3230 kJ.
Calculate the value of **B**.

B = _____ kJ


(1)

- (c) Alcohols are used as fuels.

A group of students investigated the amount of energy released when different alcohols are burned.


The students used the apparatus shown in **Figure 2**.

Figure 2



- (i) **Figure 3** shows the start temperature and the final temperature of the water.

Figure 3



Write the start temperature and the final temperature of the water in **Table 1**. Work out the increase in temperature to complete **Table 1**.

Table 1

Start temperature of the water in °C	
Final temperature of the water in °C	
Increase in temperature in °C	

(3)

- (ii) The students worked out the heat energy released by burning 1 g of each alcohol. The students used the equation:

$$\text{Heat energy released} = m \times 4.2 \times \text{increase in temperature}$$

Look at **Figure 2**. What is the value of m ?

$$m = \text{_____ g}$$

(1)

- (iii) **Table 2** shows the students' results.

Table 2

Name of alcohol	Number of carbon atoms in one molecule of alcohol	Heat energy released when 1 g of alcohol is burned in kJ
Methanol	1	11.4
Ethanol	2	13.5
Propanol	3	20.1
Butanol	4	16.8
Pentanol	5	17.2

Which value of heat energy released is anomalous?

(1)

- (iv) Look at **Table 2**.
What is the relationship between the number of carbon atoms in one molecule of alcohol and the heat energy released when 1 g of the alcohol is burned?

(1)

- (v) The value in a data book for the amount of heat energy released when 1 g of butanol is burned completely is 36.2 kJ.


Suggest two reasons why the students' result for butanol is lower than the data book value.

1. _____

2. _____

(2)

- (vi) The displayed structure of butanol is:



What is the functional group of the alcohol?

Tick (✓) **one** box.

— C — C

— C — H

— O — H


(1)

(Total 14 marks)

Q8.

This question is about energy changes in chemical reactions.


- (a) Balance the chemical equation for the combustion of methane.



(1)

(b) Alcohols are used as fuels.

A group of students investigated the amount of energy released when an alcohol was burned. The students used the apparatus shown in the diagram below.



In one experiment the temperature of 50 g of water increased from 22.0 °C to 38.4 °C.

The mass of alcohol burned was 0.8 g.

Calculate the heat energy (Q) in joules, released by burning 0.8 g of the alcohol. Use the equation:


$$Q = m \times c \times \Delta T$$

Specific heat capacity (c) = 4.2 J / g / °C


Heat energy (Q) = _____ J

(3)

(c) The chemical equation for the combustion of ethanol is:



(i) The equation for the reaction can be shown as:



Bond	Bond energy in kJ per mole
C — H	413
C — C	347

C — O	358
C = O	799
O — H	467
O = O	495

Use the bond energies to calculate the overall energy change for this reaction.

Overall energy change = _____ kJ per mole

(3)


- (ii) The reaction is exothermic.
Explain why, in terms of bonds broken and bonds formed.

(2)

- (iii) Complete the energy level diagram for the combustion of ethanol.

On the completed diagram, label:

- activation energy
- overall energy change.



(3)

(Total 12 marks)


Q9.

This question is about ethanol.

- (a) Ethanol is produced by the reaction of ethene and steam:



- (i) **Figure 1** shows the energy level diagram for the reaction.



How does the energy level diagram show that the reaction is exothermic?


(1)

- (ii) A catalyst is used for the reaction.

Explain how a catalyst increases the rate of the reaction.

(2)


- (b) **Figure 2** shows the displayed structure of ethanol.



Complete the dot and cross diagram in **Figure 3** to show the bonding in ethanol.

Show the outer shell electrons only.

Figure 3




(2)

- (c) A student burned some ethanol.

Figure 4 shows the apparatus the student used.

Figure 4



- (i) The student recorded the temperature of the water before and after heating.

His results are shown in **Table 1**.

Table 1

Temperature before heating	20.7 °C
Temperature after heating	35.1 °C

Calculate the energy used to heat the water.

Use the equation $Q = m \times c \times \Delta T$

The specific heat capacity of water = 4.2 J / g / °C

Energy used = _____ J

(3)

- (ii) **Table 2** shows the mass of the spirit burner before the ethanol was burned and after the ethanol was burned.

Table 2

Mass of spirit burner before ethanol was burned	72.80 g
Mass of spirit burner after ethanol was burned	72.10 g

Calculate the number of moles of ethanol (C_2H_5OH) that were burned.

Relative atomic masses (A_r): H = 1; C = 12; O = 16

Number of moles burned = _____

(3)

- (iii) Calculate the energy released in joules per mole.

You should assume that all the energy from the ethanol burning was used to heat the water.

Energy = _____ J / mole

(1)

- (d) The names, structures and boiling points of ethanol and two other alcohols are shown in **Table 3**.

Table 3

Name	Methanol	Ethanol	Propanol
Structure	$\begin{array}{c} \text{H} \\ \\ \text{H}-\text{C}-\text{O}-\text{H} \\ \\ \text{H} \end{array}$	$\begin{array}{c} \text{H} \quad \text{H} \\ \quad \\ \text{H}-\text{C}-\text{C}-\text{O}-\text{H} \\ \quad \\ \text{H} \quad \text{H} \end{array}$	$\begin{array}{c} \text{H} \quad \text{H} \quad \text{H} \\ \quad \quad \\ \text{H}-\text{C}-\text{C}-\text{C}-\text{O}-\text{H} \\ \quad \quad \\ \text{H} \quad \text{H} \quad \text{H} \end{array}$
Boiling point in $^{\circ}\text{C}$	65	78	97

Use your knowledge of structure and bonding to suggest why the boiling points increase as the number of carbon atoms increases.

(3)
(Total 15 marks)

Q10.

Dilute nitric acid reacts with potassium hydroxide solution.

The equation for the reaction is:



A student investigated the temperature change in this reaction.


This is the method the student used.

- Step 1 Put 25 cm³ of dilute nitric acid in a polystyrene cup.
- Step 2 Use a thermometer to measure the temperature of the dilute nitric acid.
- Step 3 Use a burette to add 4 cm³ of potassium hydroxide solution to the dilute nitric acid and stir the mixture.
- Step 4 Use a thermometer to measure the highest temperature of the mixture.
- Step 5 Repeat steps 3 and 4 until 40 cm³ of potassium hydroxide solution have been added.

The dilute nitric acid and the potassium hydroxide solution were both at room temperature.

- (a) **Figure 1** shows part of the thermometer after some potassium hydroxide solution had been added to the dilute nitric acid.

Figure 1



What is the temperature shown on the thermometer?

The temperature shown is _____ °C

(1)

(b) Errors are possible in this experiment.

(i) Suggest **two** causes of random error in the experiment.

(2)

(ii) Another student used a glass beaker instead of a polystyrene cup.


This caused a systematic error.

Why does using a glass beaker instead of a polystyrene cup cause a systematic error?

(1)

(c) The results of the student using the polystyrene cup are shown in **Figure 2**.

Figure 2



(i) How do the results in **Figure 2** show that the reaction between dilute nitric acid and potassium hydroxide solution is exothermic?

(1)

- (ii) Explain why the temperature readings decrease between 28 cm³ and 40 cm³ of potassium hydroxide solution added.

(2)

- (iii) It is difficult to use the data in **Figure 2** to find the exact volume of potassium hydroxide solution that would give the maximum temperature.

Suggest further experimental work that the student should do to make it easier to find the exact volume of potassium hydroxide solution that would give the maximum temperature

(2)

- (d) The student did further experimental work and found that 31.0 cm³ of potassium hydroxide solution neutralised 25.0 cm³ of dilute nitric acid.

The concentration of the dilute nitric acid was 2.0 moles per dm³.



Calculate the concentration of the potassium hydroxide solution in moles per dm³.

Concentration = _____ moles per dm³

(3)

- (e) The student repeated the original experiment using 25 cm³ of dilute nitric acid in a polystyrene cup and potassium hydroxide solution that was twice the original concentration.

She found that:

- a smaller volume of potassium hydroxide solution was required to reach the maximum temperature
- the maximum temperature recorded was higher.


Explain why the maximum temperature recorded was higher.

(2)
(Total 14 marks)

Q11.

Methane (CH₄) is used as a fuel.

(a) The displayed structure of methane is:



Draw a ring around a part of the displayed structure that represents a covalent bond.

(1)

(b) Why is methane a compound?

Tick (✓) **one** box.

Methane contains atoms of two elements, combined chemically.

Methane is not in the periodic table.

Methane is a mixture of two different elements.


(1)

(c) Methane burns in oxygen.

(i) The diagram below shows the energy level diagram for the complete combustion of methane.

Draw and label arrows on the diagram to show:

- the activation energy
- the enthalpy change, ΔH .



(2)

- (ii) Complete and balance the symbol equation for the complete combustion of methane.



(2)

- (iii) Explain why the **incomplete** combustion of methane is dangerous.


(2)

- (iv) Explain why, in terms of the energy involved in bond breaking and bond making, the combustion of methane is exothermic.

(3)

- (d) Methane reacts with chlorine in the presence of sunlight.

The equation for this reaction is:




Some bond dissociation energies are given in the table.

Bond	Bond dissociation energy in kJ per mole
C-H	413
C-Cl	327
Cl-Cl	243
H-Cl	432

- (i) Show that the enthalpy change, ΔH , for this reaction is -103 kJ per mole.

(3)

- (ii) Methane also reacts with bromine in the presence of sunlight.



This reaction is less exothermic than the reaction between methane and chlorine.

The enthalpy change, ΔH , is -45 kJ per mole.

What is a possible reason for this?

Tick (✓) **one** box.

CH₃Br has a lower boiling point than CH₃Cl

The C-Br bond is weaker than the C-Cl bond.

The H-Cl bond is weaker than the H-Br bond.

Chlorine is more reactive than bromine.




(1)
(Total 15 marks)

Q12.

A student investigated the temperature change when zinc reacts with copper sulfate solution.

The student used a different concentration of copper sulfate solution for each experiment.


The student used the apparatus shown below.



The student:

- measured 50 cm³ copper sulfate solution into a glass beaker
- measured the temperature of the copper sulfate solution
- added 2.3 g zinc
- measured the highest temperature
- repeated the experiment using copper sulfate solution with different concentrations.

The equation for the reaction is:



(a) The thermometer reading changes during the reaction.

Give **one** other change the student could **see** during the reaction.

(1)

(b) Suggest **one** improvement the student could make to the apparatus.

Give a reason why this improves the investigation.

Improvement _____

Reason _____

- (c) In this question you will be assessed on using good English, organising information clearly and using specialist terms where appropriate.

The student's results are shown in the table.


Table

Experiment number	Concentration of copper sulfate in moles per dm ³	Increase in temperature in °C
1	0.1	5
2	0.2	10
3	0.3	12
4	0.4	20
5	0.5	25
6	0.6	30
7	0.7	35
8	0.8	35
9	0.9	35
10	1.0	35

Describe **and** explain the trends shown in the student's results.

Q13.

A student investigates the energy released when zinc powder reacts with copper sulfate solution. The student uses the apparatus shown in **Figure 1**.



The student:

- measures 100 cm³ copper sulfate solution into a beaker
- measures the temperature of the copper sulfate solution
- puts 1 g zinc powder into the beaker
- stirs the mixture with a thermometer
- measures the highest temperature.

The student's results were:

Starting temperature = 21 °C

Highest temperature = 32 °C

- (a) (i) Calculate the change in temperature.

Change in temperature = _____ °C

(1)

- (ii) Calculate the energy released in the reaction.

Use the equation

$$\begin{array}{l} \text{energy} \\ \text{released} \\ \text{in J} \end{array} = \begin{array}{l} \text{volume of} \\ \text{solution} \\ \text{in cm}^3 \end{array} \times 4.2 \times \begin{array}{l} \text{temperature change} \\ \text{in } ^\circ\text{C} \end{array}$$

Energy released = _____ J


(2)

- (b) The reaction of zinc with copper sulfate is exothermic.

How can you tell from the student's results that the reaction is exothermic?

(1)

(c) The energy diagram for the reaction is shown in **Figure 2**.



(i) How can you tell from the energy diagram that the reaction is exothermic?

(1)

(ii) Which arrow shows the activation energy in **Figure 2**?

Tick (✓) **one** box.

A

B

C

(1)

(Total 6 marks)

Q14.

This question is about reversible reactions and chemical equilibrium.

(a) Reversible reactions can reach equilibrium in a closed system.


(i) What is meant by a closed system?

(1)

(ii) Explain why, when a reversible reaction reaches equilibrium, the reaction appears to have stopped.

(2)

- (b) In the Haber process, the reaction of nitrogen with hydrogen to produce ammonia is reversible.



- (i) Name a natural resource from which hydrogen is produced.

(1)

- (ii) The Haber process uses a catalyst to speed up the reaction.

Explain how a catalyst speeds up a reaction.


(2)

- (iii) What happens to the amount of ammonia produced at equilibrium if the pressure is increased?

Give a reason for your answer.


(2)

- (c) The decomposition of hydrogen iodide into hydrogen and iodine is reversible.



The forward reaction is endothermic.

The energy level diagram shown below is for the forward reaction.



(i) Draw an arrow to show the activation energy on the diagram.

(1)

(ii) How does the diagram show that the reaction is endothermic?

(1)

(iii) Suggest what effect, if any, increasing the temperature will have on the amount of hydrogen iodide at equilibrium.

Give a reason for your answer.

(2)


(Total 12 marks)

Q15.

A student investigates the energy released when hydrochloric acid completely neutralises sodium hydroxide solution.

The student uses the apparatus shown in **Figure 1**.

Figure 1



The student:


- measures 25 cm³ sodium hydroxide solution into a polystyrene cup
- fills a burette with hydrochloric acid
- measures the temperature of the sodium hydroxide solution
- adds 5 cm³ hydrochloric acid to the sodium hydroxide solution in the polystyrene cup
- stirs the mixture and measures the highest temperature of the mixture
- continues to add 5 cm³ portions of hydrochloric acid, stirring and measuring the highest temperature of the mixture after each addition.

(a) The student has plotted a graph of the results.

The graph line has been incorrectly drawn by including an anomalous result.

The graph is shown in **Figure 2**.

Figure 2



- (i) Suggest a cause for the anomalous result when 20 cm³ of hydrochloric acid is added.

(1)

- (ii) Suggest the true value of the temperature of the anomalous point.

Temperature = _____ °C

(1)

- (iii) What was the **total** volume of the mixture when the maximum temperature was reached?

Total volume of the mixture = _____ cm³

(1)

- (iv) Calculate the overall temperature increase in this experiment.

Overall temperature increase = _____ °C

(1)

- (v) Use your answers to (iii) and (iv) and the equation to calculate the energy released in the reaction. Give the unit.

Assume the volume in cm³ is equivalent to the mass of solution in grams.

Equation: $Q = mc\Delta T$

where:

Q = energy released
m = mass of solution (g)
c = 4.2 (J per g per °C)
 ΔT = change in temperature (°C)

Energy released = _____ Unit = _____

(2)

- (b) The student did the experiment again, starting with 50 cm³ of sodium hydroxide solution instead of 25 cm³.

Explain why this would make no difference to the overall temperature increase.


(2)

(Total 8 marks)

Q16.

Hydrogen gas is produced by the reaction of methane and steam.

- (a) The diagram represents a molecule of hydrogen.



- (i) What type of bond joins the atoms of hydrogen?

Tick (✓) **one** box.

Covalent

Metallic

Ionic

(1)

- (ii) A catalyst is used in the reaction.

Draw a ring around the correct answer to complete the sentence.

A catalyst

increases the rate of reaction.
increases the temperature.
increases the yield of a reaction.

(1)

(b) The equation for the reaction of methane and steam is:



(i) What is meant by the symbol \rightleftharpoons ?


(1)

(ii) Lowering the pressure reduces the rate of reaction.

Explain why, in terms of particles.

(2)

(iii) The graph shows the yield of hydrogen at different temperatures.



The forward reaction is endothermic.


How does the graph show that the forward reaction is endothermic?

(1)

(6)
(Total 13 marks)

Q17.

The equation for the reaction of ethene and bromine is:




The reaction is exothermic.

(a) Complete the energy level diagram.


You should label:

- the activation energy
- the enthalpy change (ΔH).



(3)

(b) (i) The equation for the reaction can be represented as:



Bond	Bond dissociation energy in kJ per mole
C—H	413
C = C	614
Br—Br	193

C—C	348
C—Br	276

Use the bond dissociation energies in the table to calculate the enthalpy change (ΔH) for this reaction.

Enthalpy change (ΔH) = _____ kJ per mole

(3)

(ii) The reaction is exothermic.

Explain why, in terms of bonds broken and bonds formed.


(2)

(Total 8 marks)

Q18.

A company manufactures ethanol (C_2H_5OH).

The reaction for the process is:



The temperature and pressure can be changed to increase the yield of ethanol at equilibrium.

(a) Explain what is meant by equilibrium.

(3)

- (b) (i) How would increasing the temperature change the **yield** of ethanol at equilibrium?

Give a reason for your answer.

(2)

- (ii) How would increasing the pressure change the **yield** of ethanol at equilibrium?

Give a reason for your answer.

(2)

- (c) A catalyst is added to increase the rate of the reaction.

Explain how adding a catalyst increases the rate of a chemical reaction.

(2)

(Total 9 marks)

Q19.


A student investigated displacement reactions of metals.

The student added different metals to copper sulfate solution and measured the temperature change.

The more reactive the metal is compared with copper, the bigger the temperature change.

The apparatus the student used is shown in **Figure 1**.

Figure 1



(a) State **three** variables that the student must control to make his investigation a fair test.

- 1. _____
- 2. _____
- 3. _____



(3)

(b) **Figure 2** shows the thermometer in one experiment before and after the student added a metal to the copper sulfate solution.

Figure 2

Before adding metal

After adding metal



Use **Figure 2** to complete **Table 1**.

Table 1

Temperature before adding metal in °C	_____
Temperature after adding metal in °C	_____
Change in temperature in °C	_____

(c) The student repeated the experiment three times with each metal.


Table 2 shows the mean temperature change for each metal.

Table 2

Metal	Mean temperature change in °C
Cobalt	4.5
Gold	0.0
Magnesium	10.0
Nickel	3.0
Silver	0.0
Tin	1.5

(i) On **Figure 3**, draw a bar chart to show the results.

Figure 3



(3)

(ii) Why is a line graph **not** a suitable way of showing the results?

(1)

(iii) Use the results to work out which metal is the most reactive.

Give a reason for your answer.

Most reactive metal _____

Reason _____

(2)

- (iv) Explain why there was no temperature change when silver metal was added to the copper sulfate solution.

(2)

- (v) It is **not** possible to put all six metals in order of reactivity using these results.

Suggest how you could change the experiment to be able to put all six metals into order of reactivity.


(2)

(Total 16 marks)

Q20.

Some students investigated reactions to produce magnesium.

- (a) The students used electrolysis to produce magnesium from magnesium chloride, as shown in the figure below.



- (i) Magnesium chloride contains magnesium ions and chloride ions.

Why does solid magnesium chloride **not** conduct electricity?

(1)

- (ii) One of the products of the electrolysis of molten magnesium chloride is magnesium.

Name the other product.

(1)

- (iii) Why do magnesium ions (Mg^{2+}) move to the negative electrode?

(1)

- (iv) At the negative electrode, the magnesium ions (Mg^{2+}) gain electrons to become magnesium atoms.

How many electrons does each magnesium ion gain?

(1)

- (b) The students did the experiment four times and weighed the magnesium produced.

The table below shows their results.

Experiment	Mass of magnesium produced in grams
1	1.13
2	0.63
3	1.11
4	1.09

- (i) There is an anomalous result.

Suggest **one** possible reason for the anomalous result.

(1)

- (ii) Calculate the mean mass of magnesium produced, taking account of the anomalous result.

Mean mass = _____ g

(2)

(c) The formula of magnesium chloride is MgCl_2

The relative formula mass of magnesium chloride is 95.

The relative atomic mass of magnesium is 24.

(i) Use the equation to calculate the percentage mass of magnesium in magnesium chloride.

$$\text{Percentage mass of magnesium} = \frac{\text{mass of magnesium}}{\text{mass of magnesium chloride}} \times 100\%$$

Percentage mass of magnesium in magnesium chloride = _____ %

(2)

(ii) Draw a ring around the relative mass of chlorine in MgCl_2

71

95

119

(1)

(d) Magnesium is also produced from the reaction of magnesium oxide with silicon.

(i) The equation for the reaction is:



What is the meaning of this symbol \rightleftharpoons ?

Draw a ring around the correct answer.

neutralisation reaction

precipitation reaction

reversible reaction

(1)

(ii) The forward reaction is endothermic.

Draw a ring around the correct answer to complete the sentence.

In an endothermic reaction the temperature of the surroundings

decreases.

increases.


stays the same.

(1)

Q21.

This question is about zinc and magnesium.

Zinc is produced by electrolysis of molten zinc chloride, as shown in the figure below.



(a) (i) Why must the zinc chloride be molten for electrolysis?

(1)

(ii) Describe what happens at the negative electrode.

(3)

(iii) Complete the half equation for the reaction at the positive electrode.



(1)

(b) Magnesium can be produced from magnesium oxide.

The equation for the reaction is:



(i) How can you tell from the equation that the reaction is done at a high temperature?

(1)

(ii) This reaction to produce magnesium from magnesium oxide is **endothermic**.

What is meant by an **endothermic** reaction?

(1)

(iii) A company made magnesium using this reaction.

Calculate the mass of magnesium oxide needed to produce 1.2 tonnes of magnesium.

Relative atomic masses (A_r): O = 16; Mg = 24

Mass of magnesium oxide needed = _____ tonnes

(3)

(iv) The company calculated that they would produce 1.2 tonnes of magnesium, but only 0.9 tonnes was produced.

Calculate the percentage yield.

Percentage yield = _____ %

(1)

(v) Give **one** reason why the calculated yield of magnesium might not be obtained.

(1)

(Total 12 marks)

Q22.

Some cars are powered by hydrogen fuel cells.

Figure 1



© Robert Couse-Baker (CC BY-SA 2.0) via Flickr

(a) What type of energy is released by hydrogen fuel cells?

Draw a ring around the correct answer.

chemical

electrical


light

(1)

(b) Owners of cars powered by fuel cells buy hydrogen from hydrogen filling stations.

Figure 2 shows how the number of hydrogen filling stations in the UK is expected to increase up to the year 2030.

Figure 2



(i) Suggest the total number of hydrogen filling stations expected in 2028.

(1)

(ii) The number of hydrogen filling stations will still be very low compared with the number of petrol filling stations.


Suggest **one** reason why.

(1)

(c) Hydrogen reacts with oxygen to produce water.

The energy level diagram for this reaction is shown in **Figure 3**.

Figure 3



Mark clearly with a cross (x) on **Figure 3** where bond breaking happens.

(1)

(Total 4 marks)

Q23.


In 1909 Fritz Haber invented a process to produce ammonia from nitrogen and hydrogen.

- (a) Complete and balance the chemical equation for the production of ammonia from nitrogen and hydrogen.



(2)

- (b) The figure below shows how the equilibrium yield of ammonia changes with pressure at different temperatures.



(i) Use the information in given in the figure to complete the sentence.
The temperature on the graph that gives the highest yield of ammonia is _____ °C.

(1)

(ii) The temperature used in the Haber process for the production of ammonia is 450 °C.

Why is a temperature much lower than 450 °C **not** used for the Haber process?

(1)

(iii) Use the information in the figure to answer this question.

Draw a ring around the pressure that gives the highest yield of ammonia.

100 200 300 400

(1)

(iv) The pressure used in the Haber process for the production of ammonia is 200 atmospheres.

Why is a pressure lower than 200 atmospheres **not** used for the Haber process?

(1)

(c) Explain how ammonia is separated from unreacted nitrogen and hydrogen in the Haber process.

(2)

(Total 8 marks)

Q24.

Some cars are powered by hydrogen fuel cells.

Figure 1



© Robert Couse-Baker (CC BY-SA 2.0) via Flickr


- (a) What type of energy is released by hydrogen fuel cells?

(1)

- (b) Owners of cars powered by fuel cells buy hydrogen from hydrogen filling stations.

Figure 2 shows how the number of hydrogen filling stations in the UK is expected to increase up to the year 2030.

Figure 2




Use the information in **Figure 2** and your own knowledge to answer this question.

Suggest **two** reasons why the UK government might encourage the building of more hydrogen filling stations.

(2)

- (c) The equation for the reaction of hydrogen with oxygen is:



During the reaction, energy is used to break the bonds of the reactants.


Energy is released when new bonds are made to form the product.

Bond energies for the reaction are given in the table below.

Bond	Bond energy in kJ
H—H	436
O=O	498
O—H	464

The structures of the reactants and product are shown in **Figure 3**.

Figure 3



- (i) Calculate the energy change for the reaction:



Energy change = _____ kJ


(3)

- (ii) The reaction of hydrogen with oxygen is exothermic.

Complete the energy level diagram for this reaction on **Figure 4**.

Clearly label the activation energy.

Figure 4



(3)
(Total 9 marks)

Q25.

Kelp is a seaweed.

Kelp can be burned to give out energy.



© Ethan Daniels/Shutterstock

- (a) Draw a ring around the correct answer to complete each sentence.

Reactions which give out energy are

- | |
|--------------|
| endothermic. |
| exothermic. |
| reversible. |

(1)

- (b) Which **two** of the following questions **cannot** be answered by scientific experiments alone?

Tick (✓) **two** boxes.

Question	Tick (✓)

How much carbon dioxide is produced when 100 g of kelp is burned?	
Does kelp give out more heat energy than coal when burned?	
Should people use kelp instead of oil as an energy source?	
Will kelp be more popular than coal in the next 10 years?	

(2)

(c) Potassium iodide can be produced from kelp.

(i) Potassium can be reacted with iodine to produce potassium iodide.



The diagram shows how this happens.

Only the outer electrons are shown.

The dots (●) and crosses (x) are used to represent electrons




Use the diagram to help you answer this question.

Describe, as fully as you can, what happens when potassium reacts with iodine to produce potassium iodide.

To get full marks you should use the words atom, electron and ion in your answer.

(4)

(ii) Potassium iodide reacts with lead nitrate.



Why is this reaction a precipitation?

(1)

(iii) How can the precipitate be removed from the reaction mixture?

(1)


(Total 9 marks)

Q26.

This question is about compounds of copper.

(a) A student made some copper(II) sulfate crystals.

The flow diagram shows the stages of the preparation of copper(II) sulfate crystals.




(i) The reaction mixture is heated in **Stage 1**.

Suggest why.

(1)

(ii) Complete the equation for this reaction.



(2)

(iii) How would the student remove the unreacted copper(II) oxide in **Stage 2**?

(1)

(iv) How would the student obtain copper(II) sulfate crystals from the copper(II) sulfate solution in **Stage 3**?

(1)


(v) The mass of crystals obtained was less than the student had calculated.

Suggest **one** reason why.

(1)

- (b) The student heated the blue copper(II) sulfate crystals.

The word equation for the reaction is shown below.



- (i) What does the symbol \rightleftharpoons mean ?

(1)

- (ii) 300 J of energy are taken in when some blue copper(II) sulfate crystals are heated.

What is the energy change when an excess of water is added to the anhydrous copper(II) sulfate produced?

(2)

- (c) A sample of copper nitride contains 3.81 g of copper and 0.28 g of nitrogen.

Calculate the empirical formula.

You **must** show all your working to get full marks.

Relative atomic masses (A_r): N = 14; Cu = 63.5.

Empirical formula = _____

(4)

(Total 13 marks)

Kelp is a seaweed.

Kelp can be used in foods and as a renewable energy source.



© Ethan Daniels/Shutterstock

- (a) Scientific experiments, on their own, **cannot** fully answer one of the following questions. Which one?

Tick (✓) **one** box.

Questions	Tick (✓)
How much carbon dioxide is produced when 100 g of kelp is burned?	
Does kelp give out more heat energy than coal?	
Will kelp last longer than coal as an energy source?	
Which fuel, kelp or coal, produces the most ash when burned?	

(1)

- (b) Scientists cannot answer the question 'should people use kelp instead of coal as an energy source?'

Give **two** reasons why.

(2)

- (c) Sodium iodide can be produced from kelp.

- (i) How many electrons are in the outer shell of an iodine atom?

(1)


- (ii) Sodium iodide contains sodium ions (Na^+) and iodide ions (I^-).

Describe, as fully as you can, what happens when sodium atoms react with iodine atoms to produce sodium iodide.

You may use a diagram in your answer

(3)

- (iii) The diagram shows the structure of sodium iodide.




Solid sodium iodide does not conduct electricity.

Why does sodium iodide solution conduct electricity?

(1)

- (iv) When sodium iodide solution is electrolysed, iodine is formed at the positive electrode.

Complete and balance the half equation for the formation of iodine.



(v) What is formed at the negative electrode when sodium iodide solution is electrolysed?


Explain why.

(2)
(Total 11 marks)

Q28.


The following steps show how to use a type of glue.

Step 1 Measure out equal amounts of the liquids from tubes **A** and **B**.




Step 2 Mix the liquids to make the glue.

Put a thin layer of the glue onto each of the surfaces to be joined.



Step 3 Put the pieces together and hold them with tape.



Step 4 Leave the glue to set.

(a) When liquids **A** and **B** are mixed a chemical reaction takes place.

This reaction is *exothermic*.

What does *exothermic* mean?

(2)

- (b) The time taken for the glue to set at different temperatures is given in the table below.

Temperature in °C	Time taken for the glue to set
20	3 days
60	6 hours
90	1 hour

- (i) Use the correct answer from the box to complete each sentence.

decreases	increases	stays the same
------------------	------------------	-----------------------

When the temperature is increased the time taken for the glue to set

When the temperature is increased the rate of the setting reaction

(2)

- (ii) Tick (✓) **two** reasons why an increase in temperature affects the rate of reaction.

Reason	Tick (✓)
It gives the particles more energy	
It increases the concentration of the particles	
It increases the surface area of the particles	
It makes the particles move faster	

(2)


(Total 6 marks)

Q29.


Read the information about energy changes and then answer the questions.

A student did an experiment to find the energy change when hydrochloric acid reacts with sodium hydroxide.

The equation which represents the reaction is:



The student used the apparatus shown in the diagram.



The student placed 50 cm³ of hydrochloric acid in a glass beaker and measured the initial temperature.

The student then quickly added 50 cm³ of sodium hydroxide solution and stirred the mixture with the thermometer. The highest temperature was recorded.

The student repeated the experiment, and calculated the temperature change each time.

	Experiment 1	Experiment 2	Experiment 3	Experiment 4
Initial temperature in °C	19.0	22.0	19.2	19.0
Highest temperature in °C	26.2	29.0	26.0	23.5
Temperature change in °C	7.2	7.0	6.8	4.5

- (a) The biggest error in this experiment is heat loss.

Suggest how the apparatus could be modified to reduce heat loss.

(1)

- (b) Suggest why it is important to mix the chemicals thoroughly.

(1)

- (c) Which **one** of these experiments was probably done on a different day to the others?

Give a reason for your answer.

(1)

- (d) Suggest why experiment **4** should **not** be used to calculate the average temperature change.

(1)

- (e) Calculate the average temperature change from the first three experiments.

Answer = _____ °C

(1)

- (f) Use the following equation to calculate the energy change for this reaction.


$$\text{Energy change in joules} = 100 \times 4.2 \times \text{average temperature change}$$

Answer = _____ J

(1)

- (g) Which **one** of these energy level diagrams represents the energy change for this reaction?

Give a reason for your answer.



(1)

(Total 7 marks)


Q30.

Methanol (CH₃OH) can be made by reacting methane (CH₄) and oxygen (O₂). The reaction is exothermic.

The equation for the reaction is:



- (a) The energy level diagram for this reaction is given below.




- (i) How does the diagram show that this reaction is exothermic?

(1)

- (ii) A platinum catalyst can be used to increase the rate of this reaction.
What effect does adding a catalyst have on the energy level diagram?

(1)

- (b) The equation can also be written showing the structural formulae of the reactants and the product.



- (i) Use the bond energies given in the table to help you to calculate the energy change for this reaction.

Bond	Bond energy in kJ
C—H	435
O=O	497
C—O	336

O—H	464
-----	-----

Energy change = _____ kJ

(3)


(iii) In terms of the bond energies, why is this an exothermic reaction?

(1)

(Total 6 marks)

Q31.

The diagram shows one way of producing iron.



Iron oxide reacts with aluminium to produce iron.

The symbol equation for the reaction is:



(a) (i) Complete the word equation for this reaction.

iron oxide + aluminium \longrightarrow iron + _____

(1)

(ii) The magnesium ribbon is lit to start the reaction.

Why does the burning magnesium ribbon start the reaction?


(1)

- (b) In industry, iron is produced in the blast furnace when iron oxide is heated with carbon.

The iron from the blast furnace is called cast iron.

Cast iron contains carbon.

The diagrams show the structure of pure iron and cast iron.



Use the diagrams to help you answer the questions.

- (i) Draw a ring around the correct answer to complete the sentence.

Pure iron is an element because pure iron


contains only one sort of atom.
is magnetic.
is a metal.

(1)

- (ii) Suggest why cast iron is harder than pure iron.

(2)

- (c) Aluminium is extracted by electrolysis using the ionic compound aluminium oxide.



(i) Aluminium **cannot** be extracted by heating aluminium oxide with carbon.

Suggest why.


(1)

(ii) Why is aluminium oxide dissolved in molten cryolite?

(1)

(iii) Aluminium metal is produced at the negative electrode (cathode).

Complete the half equation for the process.



(1)

(iv) Use the half equation to state why Al^{3+} ions are reduced.

(1)

(v) Explain why the positive electrodes (anodes) burn away.

Use your knowledge of the products of electrolysis to help you.

(4)

(Total 13 marks)

Q32.

(a) Which sub-atomic particles are present in the nucleus of an atom?

_____ and _____

(2)

- (b) There are two isotopes of the element chlorine:



Describe, in terms of sub-atomic particles, **one** similarity and **one** difference between atoms of the two isotopes of chlorine.

Similarity _____

Difference _____


(2)

- (c) Chlorine reacts with hydrogen to produce hydrogen chloride.

- (i) The table shows the values of some bond dissociation energies.

Bond	H—H	Cl—Cl	H—Cl
Dissociation energy in kJ per mole	436	242	431

Use the values in the table to calculate the enthalpy change (ΔH) for the reaction.



Enthalpy change (ΔH) = _____ kJ per mole

(3)

- (ii) Hydrogen also reacts with fluorine.



Draw an energy level diagram for this reaction.

Include on your diagram labels to show:

- the reactants and the products
- the overall enthalpy change (ΔH)

- the activation energy.

(3)
(Total 10 marks)

Q33.

When ammonium chloride is dissolved in water, there is a temperature change.

A student investigated how the temperature of water changed when different masses of ammonium chloride were added to the same volume of water.

The water used was at room temperature.

The student's results are shown in the table.

Mass of ammonium chloride in g	Final temperature of solution in °C
10	14.5
20	8.5
25	5.5
30	2.5
35	1.0
40	1.0
45	1.0

- (a) (i) Use the correct word from the box to complete the sentence.

endothermic	exothermic	reduction
--------------------	-------------------	------------------

When ammonium chloride dissolves in water, the change can be described as _____.

(1)

- (ii) Give a reason for your answer to part (a) (i). Refer to the table of results in your answer.

(1)

- (b) The student added the ammonium chloride to water and stirred the mixture.

The water was in a glass beaker.

His teacher said that using a glass beaker could cause inaccurate results.

What could the student have used instead of a glass beaker to improve the accuracy?

Give a reason why this would improve the accuracy of his results.

(2)

- (c) The student made sure his investigation was a fair test.

State **two** control variables the student should keep the same.

Give a reason why changing each of these two control variables would affect the temperature change.

Control variable 1 _____

Reason _____

Control variable 2 _____

Reason _____


(4)

- (d) (i) The student's results table has been repeated below.

Mass of ammonium	Final temperature
------------------	-------------------

chloride in g	of solution in °C
10	14.5
20	8.5
25	5.5
30	2.5
35	1.0
40	1.0
45	1.0

Plot the results on the grid.



(2)

(ii) Complete the graph by drawing two straight lines of best fit through the points.

(2)

(iii) Use the graph to estimate the temperature of the room.

Show your working on the graph.

Temperature of room = _____ °C

(2)

- (e) Explain why the final temperature was the same for all masses of 35 g and greater.

(2)

- (f) A second student also did one of the experiments.

This student recorded a final temperature of 14.5 °C.

Both students dissolved 20 g of ammonium chloride in water.


Use the graph to explain the difference in the two final temperatures.

(2)

(Total 18 marks)

Q34.

- (a) Some students did an experiment to find the temperature change when hydrochloric acid reacts with sodium hydrogencarbonate.



The results are in the table.

Number of spatula measures of sodium hydrogencarbonate	Start temperature in °C	Final temperature in °C	Change in temperature in °C
2	20	16	4
4	20	14	6

6	19	11	8
8	20	10	10
10	19	9	10
12	20	10	10

- (i) Describe, as fully as you can, the trends shown in the students' results.

(3)


- (ii) State the type of energy transfer for this reaction.

(1)

- (b) Sodium hydrogencarbonate is used as baking powder for making cakes.

When the cake mixture is baked the sodium hydrogencarbonate decomposes.

The equation for the reaction is:



- (i) The cake mixture rises when baked.




© Michael Valdez/iStock

Use the equation to suggest why.

(1)

- (ii) The same reaction can be reversed to produce sodium hydrogencarbonate from sodium carbonate.



Do the reactants need to be heated?

Give a reason for your answer.

(1)

- (c) (i) Calculate the relative formula mass of sodium hydrogencarbonate (NaHCO_3).

Relative atomic masses (A_r): H=1; C=12; O=16; Na=23

Relative formula mass (M_r) = _____

(2)


- (ii) Calculate the percentage by mass of carbon in sodium hydrogencarbonate.

Percentage of carbon = _____ %

(1)

Q35.

A camping stove uses propane gas.



(a) A student did an experiment to find the energy released when propane is burned.

The student:

- put 500 g water into a beaker
- measured the temperature of the water
- heated the water by burning propane for 1 minute
- measured the temperature of the water again.

The student found the temperature change was 20 °C.

The student can calculate the energy released, in joules (J), using the equation:

$$\text{energy released (J)} = \text{mass of water (g)} \times 4.2 \times \text{temperature change (}^\circ\text{C)}$$

(i) Use the student's result to calculate the energy released in joules (J).

Energy released = _____ J

(2)

(ii) State **two** safety precautions that the student should take during the experiment.

1. _____

2. _____


(2)

- (iii) Tick (✓) **two** boxes which describe how the student could make his result more accurate.

	Tick (✓)
Stir the water before measuring the temperature.	
Heat the water until it boils.	
Place a lid on the beaker.	
Use a larger beaker for the water.	

(2)

- (b) The change in energy when propane is burned can be shown in an energy level diagram.



Draw **one** line from each description to the correct letter.

Description	Letter
<input type="text" value="products"/>	<input type="text" value="P"/>
<input type="text" value="activation energy"/>	<input type="text" value="Q"/>
<input type="text" value="energy released by the reaction"/>	<input type="text" value="R"/>
	<input type="text" value="S"/>

(3)

(c) Propane and hydrogen are both used as fuels.

Some information about propane and hydrogen is given in the table.

Fuel	Resource	Products formed when fuel burned
propane	crude oil	carbon dioxide and water
hydrogen	water	water

Use the information in the table to suggest **two** disadvantages that propane has as a fuel compared to hydrogen.

1. _____

2. _____

(2)
(Total 11 marks)

