


## Quantitative Chemistry


### Q1.

A student investigated the reaction of copper carbonate with dilute sulfuric acid.

The student used the apparatus shown in the figure below.



- (a) Complete the state symbols in the equation.



(2)

- (b) Why did the balance reading decrease during the reaction?

Tick **one** box.

The copper carbonate broke down.

A salt was produced in the reaction.

A gas was lost from the flask.

Water was produced in the reaction.

(1)

- (c) Describe a safe method for making pure crystals of copper sulfate from copper carbonate and dilute sulfuric acid. Use the information in the figure above to help you.

In your method you should name all of the apparatus you will use.

---

---

---

---

---

---

---

---

---

---


---

(6)

(d) The percentage atom economy for a reaction is calculated using:

$$\frac{\text{Relative formula mass of desired product from equation}}{\text{Sum of relative formula masses of all reactants from equation}} \times 100$$

The equation for the reaction of copper carbonate and sulfuric acid is:



Relative formula masses :  $\text{CuCO}_3 = 123.5$ ;  $\text{H}_2\text{SO}_4 = 98.0$ ;  $\text{CuSO}_4 = 159.5$

Calculate the percentage atom economy for making copper sulfate from copper carbonate.

---

---

---

---

---

Atom economy = \_\_\_\_\_ %

(3)

(e) Give **one** reason why is it important for the percentage atom economy of a reaction to be as high as possible.

---

---


(1)

(Total 13 marks)

### Q2.

**Figure 1** shows the outer electrons in an atom of the Group 1 element potassium and in an atom of the Group 6 element sulfur.

**Figure 1**



(a) Potassium forms an ionic compound with sulfur.

Describe what happens when **two** atoms of potassium react with **one** atom of sulfur.

Give your answer in terms of electron transfer.

Give the formulae of the ions formed.

---

---

---

---

---

---

---

---

---

---


---

---

(5)

- (b) The structure of potassium sulfide can be represented using the ball and stick model in **Figure 2**.

**Figure 2**



The ball and stick model is **not** a true representation of the structure of potassium sulfide.

Give **one** reason why.

---


---

(1)

- (c) Sulfur can also form covalent bonds.

Complete the dot and cross diagram to show the covalent bonding in a molecule of hydrogen sulfide.

Show the outer shell electrons only.



(2)

(d) Calculate the relative formula mass ( $M_r$ ) of aluminium sulfate  $\text{Al}_2(\text{SO}_4)_3$

Relative atomic masses ( $A_r$ ): oxygen = 16; aluminium = 27; sulfur = 32

---



---



---

Relative formula mass = \_\_\_\_\_

(2)

(e) Covalent compounds such as hydrogen sulfide have low melting points and do **not** conduct electricity when molten.

Draw **one** line from each property to the explanation of the property.

**Property**

**Explanation  
of property**

Low melting point	Electrons are free to move
	There are no charged particles free to move
	Ions are free to move
Does not conduct electricity when	Weak intermolecular forces of attraction

molten

Bonds are weak

Bonds are strong

(2)

- (f) Ionic compounds such as potassium sulfide have high boiling points and conduct electricity when dissolved in water.

Draw **one** line from each property to the explanation of the property.

**Property**

**Explanation  
of property**

High boiling point

Electrons are free  
to move

There are no  
charged particles  
free to move

Ions are free  
to move

Weak intermolecular  
forces of attraction

Conduct electricity  
when molten

Bonds are weak

Bonds are strong

(2)

(Total 14 marks)

**Q3.**

A student investigated the reactions of copper carbonate and copper oxide with dilute hydrochloric acid.

In both reactions one of the products is copper chloride.

- (a) Describe how a sample of copper chloride crystals could be made from copper carbonate and dilute hydrochloric acid.

---

---

---

---

---

---

---

---


---

---

(4)

- (b) A student wanted to make 11.0 g of copper chloride.

The equation for the reaction is:



Relative atomic masses,  $A_r$ : H = 1; C = 12; O = 16; Cl = 35.5; Cu = 63.5

Calculate the mass of copper carbonate the student should react with dilute hydrochloric acid to make 11.0 g of copper chloride.

---

---

---

---

---

Mass of copper carbonate = \_\_\_\_\_ g

(4)

- (c) The percentage yield of copper chloride was 79.1 %.

Calculate the mass of copper chloride the student actually produced.



---

---

Actual mass of copper chloride produced = \_\_\_\_\_ g

(2)

- (d) Look at the equations for the two reactions:



Reactive formula masses: CuO = 79.5; HCl = 36.5; CuCl<sub>2</sub> = 134.5; H<sub>2</sub>O = 18

The percentage atom economy for a reaction is calculated using:

$$\frac{\text{Relative formula mass of desired product from equation}}{\text{Sum of relative formula masses of all reactants from equation}} \times 100$$

Calculate the percentage atom economy for Reaction 2.

---

---

---

---

---

---

---

Percentage atom economy = \_\_\_\_\_ %

(3)

- (e) The atom economy for Reaction 1 is 68.45 %.  
Compare the atom economies of the two reactions for making copper chloride.

Give a reason for the difference.

---


---

(1)

(Total 14 marks)

#### Q4.

Sodium carbonate reacts with dilute hydrochloric acid:




A student investigated the volume of carbon dioxide produced when different masses of sodium carbonate were reacted with dilute hydrochloric acid.

This is the method used.

1. Place a known mass of sodium carbonate in a conical flask.
2. Measure 10 cm<sup>3</sup> of dilute hydrochloric acid using a measuring cylinder.
3. Pour the acid into the conical flask.
4. Place a bung in the flask and collect the gas until the reaction is complete.

- (a) The student set up the apparatus as shown in the figure below.



Identify the error in the way the student set up the apparatus.

Describe what would happen if the student used the apparatus shown.

---



---



---



---

(2)

(b) The student corrected the error.

The student's results are shown in the table below.

Mass of sodium carbonate in g	Volume of carbon dioxide gas in cm <sup>3</sup>
0.07	16.0
0.12	27.5
0.23	52.0
0.29	12.5
0.34	77.0
0.54	95.0
0.59	95.0
0.65	95.0

The result for 0.29 g of sodium carbonate is anomalous.

Suggest what may have happened to cause this anomalous result.

---



---

(1)

(c) Why does the volume of carbon dioxide collected stop increasing at 95.0 cm<sup>3</sup>?

---



---

(1)

- (d) What further work could the student do to be more certain about the minimum mass of sodium carbonate needed to produce 95.0 cm<sup>3</sup> of carbon dioxide?

---

---

(1)

- (e) The carbon dioxide was collected at room temperature and pressure.  
The volume of one mole of any gas at room temperature and pressure is 24.0 dm<sup>3</sup>.

How many moles of carbon dioxide is 95.0 cm<sup>3</sup>?

Give your answer in three significant figures.

---

---

---

---

\_\_\_\_\_ mol

(2)

- (f) Suggest **one** improvement that could be made to the apparatus used that would give more accurate results.

Give a reason for your answer.

---

---

---

---

(2)

- (g) One student said that the results of the experiment were wrong because the first few bubbles of gas collected were air.

A second student said this would make no difference to the results.

Explain why the second student was correct.

---

---

---


---

---

**Q5.**

Sodium hydroxide neutralises sulfuric acid.

The equation for the reaction is:



- (a) Sulfuric acid is a strong acid.

What is meant by a strong acid?

---

---

---

(2)

- (b) Write the ionic equation for this neutralisation reaction. Include state symbols.

---

(2)

- (c) A student used a pipette to add 25.0 cm<sup>3</sup> of sodium hydroxide of unknown concentration to a conical flask.

The student carried out a titration to find out the volume of 0.100 mol / dm<sup>3</sup> sulfuric acid needed to neutralise the sodium hydroxide.

Describe how the student would complete the titration.

You should name a suitable indicator and give the colour change that would be seen.

---

---

---

---

---

---

---

---

---

---

---

(4)

(d) The student carried out five titrations. Her results are shown in the table below.

	Titration 1	Titration 2	Titration 3	Titration 4	Titration 5
Volume of 0.100 mol / dm <sup>3</sup> sulfuric acid in cm <sup>3</sup>	27.40	28.15	27.05	27.15	27.15

Concordant results are within 0.10 cm<sup>3</sup> of each other.

Use the student's concordant results to work out the mean volume of 0.100 mol / dm<sup>3</sup> sulfuric acid added.

---

---


---

---

Mean volume = \_\_\_\_\_ cm<sup>3</sup>

(2)

(e) The equation for the reaction is:



Calculate the concentration of the sodium hydroxide.

Give your answer to three significant figures.

---

---

---

---

---

---

---

---

Concentration = \_\_\_\_\_ mol / dm<sup>3</sup>

(4)

(f) The student did another experiment using 20 cm<sup>3</sup> of sodium hydroxide solution with a concentration of 0.18 mol / dm<sup>3</sup>.

Relative formula mass ( $M_r$ ) of NaOH = 40

Calculate the mass of sodium hydroxide in 20 cm<sup>3</sup> of this solution.

---

---

Mass = \_\_\_\_\_ g


(2)

(Total 16 marks)

**Q6.**

- (a) Nitrogen and hydrogen are passed over iron to produce ammonia in the Haber Process.

Balance the equation for the reaction.



(1)

- (b) What is iron used for in the Haber process?

Tick **one** box.

catalyst


fuel

monomer

reactant

(1)

- (c) The figure below shows how the percentage yield of ammonia changes with pressure.



Describe the trend shown in the figure above.

---

---

(1)

- (d) Use the figure above to determine the difference in percentage yield of ammonia at 150 atmospheres pressure and 250 atmospheres pressure.

Difference in percentage yield of ammonia = \_\_\_\_\_ %


(2)

(Total 5 marks)

### Q7.

A student investigated the rate of reaction between marble chips and hydrochloric acid.

**Figure 1** shows the apparatus the student used.



- (a) What is **A**?

Tick **one** box.

cotton wool

limestone

poly(ethene)

rubber bung

(1)

- (b) **Table 1** shows the student's results for one investigation.

**Table 1**


Time in s	Mass lost in g
0	0.0
20	1.6

40	2.6
60	2.9
80	3.7
100	4.0
120	4.0

On **Figure 2**:

- Plot these results on the grid.
- Draw a line of best fit.

**Figure 2**



(3)


(c) Use **Figure 2** to complete **Table 2**.

**Table 2**

Mass lost after 0.5 minutes	_____ g
Time taken to complete the reaction	_____ s

(2)

(d) The equation for the reaction is:



Explain why there is a loss in mass in this investigation.

---



---

---

---

(2)

- (e) Another student investigated the rate of a different reaction.

**Table 3** shows the results from the different reaction.

**Table 3**

Mass lost when the reaction was complete	9.85 g
Time taken to complete the reaction	2 minutes 30 seconds

Calculate the mean rate of the reaction using **Table 3** and the equation:

$$\text{mean rate of reaction} = \frac{\text{mass lost in g}}{\text{time taken in s}}$$

Give your answer to two decimal places.

---

---

Mean rate of reaction = \_\_\_\_\_ g / s

(2)

- (f) The student measured the change in mass of the reactants.

Describe another method, other than measuring the change in mass of the reactions, that the student could have used to find the rate of the reaction between marble chips and hydrochloric acid.

---

---

---

---

(2)

- (g) Another student planned to investigate the effect of temperature on the rate of reaction.

The student predicted that the rate of reaction would increase as the temperature was increased.

Give **two** reasons why the student's prediction is correct.

Tick **two** boxes.

The particles are more concentrated.

The particles have a greater mass.

The particles have a larger surface area.

The particles have more energy.

The particles move faster.

(2)

(Total 14 marks)

**Q8.**

Metals are extracted from ores in the Earth's crust.

(a) Why is copper used in the manufacture of computers?

Tick (✓) **one** box.

Because it has a high density.

Because it does not react with water.


Because it is a good conductor of electricity.

(1)

(b) **Figure 1** shows the percentage (%) by mass of some metals in the Earth's crust.

**Figure 1**





(i) What is the percentage by mass of magnesium in the Earth's crust?

\_\_\_\_\_ %

(1)

(ii) On **Figure 1** draw the bars for:

- calcium at 3.6% by mass
- iron at 5.0% by mass.

(2)

(c) An ore of zinc contains zinc carbonate.

The equation for the reaction when zinc carbonate is heated is:



(i) What is the name of this type of reaction?

Tick (✓) **one** box.

corrosion

decomposition

electrolysis

(1)

(ii) Which substance in the equation is a gas at room temperature (20 °C)?

Tick (✓) **one** box.

zinc carbonate

zinc oxide

carbon dioxide

(1)

(iii) Complete the table below to show the number of atoms of carbon and oxygen in the formula of zinc carbonate.

Element	Number of atoms in the formula $\text{ZnCO}_3$
zinc, Zn	1
carbon, C	
oxygen, O	

(2)

(iv) When 125 g zinc carbonate is heated, 81 g zinc oxide is produced.


Calculate the mass of carbon dioxide produced.

\_\_\_\_\_g

(1)

(d) **Figure 2** shows a simple life cycle of a car body.

**Figure 2**



(i) What is **one** reason why iron from the blast furnace is converted into steel?

Tick (✓) **one** box.

To make the iron pure.

To make the iron more brittle.

To make alloys for specific uses.

(1)

(ii) Apart from cost, give **three different** reasons why steel should be recycled.

1. \_\_\_\_\_

\_\_\_\_\_

2. \_\_\_\_\_

\_\_\_\_\_

3. \_\_\_\_\_

\_\_\_\_\_

(3)

(Total 13 marks)

### Q9.

Metals are extracted from ores in the Earth's crust.

Some ores contain metal carbonates and some ores contain metal oxides.

(a) (i) Name the type of reaction that happens when a metal carbonate is heated.

\_\_\_\_\_

(1)

(ii) Which solid product is formed when copper carbonate is heated?

Tick (✓) **one** box.

copper

copper nitrate

copper oxide


copper sulfide

(1)

(b) A student investigated heating four metal carbonates.

**Figure 1** shows the apparatus used.

**Figure 1**



The student heated each metal carbonate for five minutes.

The table below shows the results.

Metal carbonate	Mass of metal carbonate at start in g	Mass of solid after heating for 5 minutes in g	Observations
Copper carbonate	10.0	6.9	Limewater turns cloudy
Magnesium carbonate	10.0	9.1	Limewater turns cloudy
Potassium carbonate	10.0	10.0	Limewater does not turn cloudy
Zinc carbonate	10.0	8.3	Limewater turns cloudy

(i) Explain the results for potassium carbonate.

---



---



---



---



---



---



---



---

(3)

(ii) Suggest how the reactivity series can be used to predict which metal carbonate reacts most easily when heated.

---



---




---

(2)

(c) **Figure 2** shows a simple life cycle of a car body.

**Figure 2**



(i) Complete the sentence.


Iron ores must contain enough iron to \_\_\_\_\_

\_\_\_\_\_

(1)

(ii) Some iron ores contain iron oxide ( $\text{Fe}_2\text{O}_3$ ).

Complete and balance the equation for a reaction to produce iron from iron oxide.



(2)

(iii) Give **two** reasons why iron produced in a blast furnace is converted into steel.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

(2)

(iv) When a car reaches the end of its useful life, the car body can be:

- recycled
- reused
- sent to landfill.

Give **three** reasons why a steel car body should be recycled and **not** reused or sent to landfill.

---

---

---

---

---

---

---


(3)  
(Total 15 marks)

**Q10.**

**In this question you will be assessed on using good English, organising information clearly and using specialist terms where appropriate.**

A student has to check if two samples of hydrochloric acid, **A** and **B**, are the same concentration.

Describe how the student could use the apparatus and the solutions in the diagram below to carry out titrations.




(Total 6 marks)

**Q11.**

The graph in **Figure 1** shows a flow diagram for the Haber process.

**Figure 1**



- (a) (i) Hydrogen gas is obtained from methane.  
Name **one** source of methane.

(1)

(ii) Air is the source used to produce nitrogen for the Haber process. Suggest why air must **not** get into the reactor.

---

---

---

(2)

(iii) Describe what happens to the mixture of gases from the reactor.

---

---

---

---


---

---

(3)

(b) The graph in **Figure 2** shows the percentage yield of ammonia using different conditions.

**Figure 2**





- (i) Use **Figure 2** to suggest the conditions that produce the greatest yield of ammonia.

---



---

(1)

- (ii) Use **Figure 2** to suggest and explain why the conditions used to produce ammonia in the Haber process are a temperature of 450 °C and a pressure of 200 atmospheres.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

(5)


(Total 12 marks)

**Q12.**

This question is about iron and aluminium.

- (a) Iron is extracted in a blast furnace. **Figure 1** is a diagram of a blast furnace.

**Figure 1**



- (i) Calcium carbonate decomposes at high temperatures.

Complete the word equation for the decomposition of calcium carbonate.



\_\_\_\_\_

(2)

- (ii) Carbon burns to produce carbon dioxide.

The carbon dioxide produced reacts with more carbon to produce carbon monoxide.

Balance the equation.



(1)

- (iii) Carbon monoxide reduces iron(III) oxide:



Calculate the maximum mass of iron that can be produced from 300 tonnes of iron(III) oxide.

Relative atomic masses ( $A_r$ ): O = 16; Fe = 56

---

---

---

---


---

---

Maximum mass = \_\_\_\_\_ tonnes

(3)

(b) Aluminium is extracted by electrolysis, as shown in **Figure 2**.



(i) Why can aluminium **not** be extracted by heating aluminium oxide with carbon?

---

---

(1)

(ii) Explain why aluminium forms at the negative electrode during electrolysis.

---

---

---

---

---

---

---

---

(3)

(iii) Explain how carbon dioxide forms at the positive electrodes during electrolysis.

---

---

---

---

---

---

---

---

(3)

(Total 13 marks)

**Q13.**

Dilute nitric acid reacts with potassium hydroxide solution.

The equation for the reaction is:




A student investigated the temperature change in this reaction.

This is the method the student used.

- Step 1 Put 25 cm<sup>3</sup> of dilute nitric acid in a polystyrene cup.
- Step 2 Use a thermometer to measure the temperature of the dilute nitric acid.
- Step 3 Use a burette to add 4 cm<sup>3</sup> of potassium hydroxide solution to the dilute nitric acid and stir the mixture.
- Step 4 Use a thermometer to measure the highest temperature of the mixture.
- Step 5 Repeat steps 3 and 4 until 40 cm<sup>3</sup> of potassium hydroxide solution have been added.

The dilute nitric acid and the potassium hydroxide solution were both at room temperature.

- (a) **Figure 1** shows part of the thermometer after some potassium hydroxide solution had been added to the dilute nitric acid.



What is the temperature shown on the thermometer?

The temperature shown is \_\_\_\_\_ °C

(1)

- (b) Errors are possible in this experiment.

- (i) Suggest **two** causes of random error in the experiment.

---

---

---

---

(2)

- (ii) Another student used a glass beaker instead of a polystyrene cup.

This caused a systematic error.

Why does using a glass beaker instead of a polystyrene cup cause a systematic error?

---


---

---

(1)

- (c) The results of the student using the polystyrene cup are shown in **Figure 2**.

**Figure 2**



- (i) How do the results in **Figure 2** show that the reaction between dilute nitric acid and potassium hydroxide solution is exothermic?

---

---

(1)

- (ii) Explain why the temperature readings decrease between 28 cm<sup>3</sup> and 40 cm<sup>3</sup> of potassium hydroxide solution added.

---

---

---

(2)

- (iii) It is difficult to use the data in **Figure 2** to find the exact volume of potassium hydroxide solution that would give the maximum temperature.

Suggest further experimental work that the student should do to make it easier to find the exact volume of potassium hydroxide solution that would give the maximum temperature

---

---

---

---

(2)

- (d) The student did further experimental work and found that 31.0 cm<sup>3</sup> of potassium hydroxide solution neutralised 25.0 cm<sup>3</sup> of dilute nitric acid.

The concentration of the dilute nitric acid was 2.0 moles per dm<sup>3</sup>.



Calculate the concentration of the potassium hydroxide solution in moles per dm<sup>3</sup>.

---

---

---

---

---

---

---

Concentration = \_\_\_\_\_ moles per dm<sup>3</sup>

(3)

- (e) The student repeated the original experiment using 25 cm<sup>3</sup> of dilute nitric acid in a polystyrene cup and potassium hydroxide solution that was twice the original concentration.

She found that:

- a smaller volume of potassium hydroxide solution was required to reach the maximum temperature
- the maximum temperature recorded was higher.

Explain why the maximum temperature recorded was higher.

---

---

---

---

(2)

(Total 14 marks)

### Q14.

Copper is a transition metal.

- (a) (i) Where is copper in the periodic table?

Tick (✓) **one** box.

in the central block

in Group 1

in the noble gas group

(1)

(ii) What is a property of copper?

Tick (✓) **one** box.

breaks easily

conducts electricity

does not conduct heat

(1)

(b) Copper ores are quarried by digging large holes in the ground, as shown in **Figure 1**.

**Figure 1**



Give **two** reasons why quarrying is bad for the environment.

---



---



---



---

(2)


(c) Some copper ores contain only 2% copper.

Most of the ore is rock that is not needed.

In one ore, the main compound is copper carbonate ( $\text{CuCO}_3$ ).

**Figure 2** shows the stages used in the extraction of copper from this ore.

**Figure 2**



(i) Why is **Stage 2** important?

---



---

(1)

(ii) The equation for the reaction in **Stage 3** is:



From the symbol equation, a company calculated that 247 tonnes of copper carbonate are needed to produce 127 tonnes of copper and 132 tonnes of carbon dioxide are released.

Calculate the mass of carbon needed to make 127 tonnes of copper.

copper carbonate	+	carbon	→	copper	+	carbon dioxide
247 tonnes		..... tonnes		127 tonnes		132 tonnes



---

---

(2)

- (iii) Suggest **one** reason why it is important for the company to calculate the mass of reactants in **Stage 3**.

---

---

(1)

(Total 8 marks)

**Q15.**

This question is about carbon and gases in the air.

- (a) Carbon atoms have protons, neutrons and electrons.

Complete the table by writing the relative mass of a neutron and an electron.

Name of particle	Relative mass
proton	1
neutron	
electron	

(2)

- (b) What is the total number of protons and neutrons in an atom called?

Tick (✓) **one** box.

The atomic number


The mass number

One mole of the atom

(1)

- (c) An atom of carbon has six electrons.

Which structure, **A**, **B** or **C**, represents the electronic structure of the carbon atom?



The carbon atom is structure

(1)

(d) Carbon reacts with oxygen to produce carbon dioxide (CO<sub>2</sub>).

(i) How many different elements are in one molecule of carbon dioxide?

\_\_\_\_\_

(1)

(ii) What is the total number of atoms in one molecule of carbon dioxide?

\_\_\_\_\_

(1)

(e) Sometimes carbon reacts with oxygen to produce carbon monoxide (CO).

(i) Calculate the relative formula mass ( $M_r$ ) of carbon monoxide.

Relative atomic masses ( $A_r$ ): C = 12; O = 16

\_\_\_\_\_  
\_\_\_\_\_

$M_r$  of carbon monoxide = \_\_\_\_\_

(1)

(ii) Calculate the percentage by mass of carbon in carbon monoxide.


\_\_\_\_\_  
\_\_\_\_\_

Percentage by mass of carbon in carbon monoxide = \_\_\_\_\_%

(1)

(f) Carbon dioxide is one of the gases in the air.

(i) The graph shows the percentage of argon and the percentage of carbon dioxide in the air.



What is the percentage of argon in the air?

Percentage of argon = \_\_\_\_\_ %

(1)

- (ii) An instrumental method is used to measure the amount of carbon dioxide in the air.

Give **one** reason for using an instrumental method.

---



---

(1)

(Total 10 marks)

### Q16.

This question is about atoms and isotopes.

- (a) Atoms contain protons, neutrons and electrons.

A lithium atom has the symbol  ${}^7_3\text{Li}$

Explain, in terms of sub-atomic particles, why the mass number of this lithium atom is 7.

---



---



---



---



---



---

(3)

- (b) Amounts of substances can be described in different ways.

Complete the sentences.

One mole of a substance is the relative formula mass in

---

The relative atomic mass of an element compares the mass of an atom of an element with the mass of an atom of

---

(2)

- (c) Two isotopes of oxygen are  $^{18}_8\text{O}$  and  $^{16}_8\text{O}$

Describe the similarities and differences between the isotopes  $^{18}_8\text{O}$  and  $^{16}_8\text{O}$

You should refer to the numbers of sub-atomic particles in each isotope.

---

---

---

---

---

---

---

(3)

(Total 8 marks)

### Q17.

This question is about chemical analysis.

- (a) A student has solutions of three compounds, X, Y and Z.

The student uses tests to identify the ions in the three compounds.

The student records the results of the tests in the table.

Compound	Test			
	Flame test	Add sodium hydroxide solution	Add hydrochloric acid and barium chloride solution	Add nitric acid and silver nitrate solution
X	no colour	green precipitate	white precipitate	no reaction
Y	yellow flame	no reaction	no reaction	yellow precipitate

<b>Z</b>	no colour	brown precipitate	no reaction	cream precipitate
----------	-----------	-------------------	-------------	-------------------

Identify the **two** ions present in each compound, **X**, **Y** and **Z**.

**X** \_\_\_\_\_

**Y** \_\_\_\_\_

**Z** \_\_\_\_\_

(3)

- (b) A chemist needs to find the concentration of a solution of barium hydroxide. Barium hydroxide solution is an alkali.

The chemist could find the concentration of the barium hydroxide solution using two different methods.

**Method 1**

- An excess of sodium sulfate solution is added to 25 cm<sup>3</sup> of the barium hydroxide solution. A precipitate of barium sulfate is formed.
- The precipitate of barium sulfate is filtered, dried and weighed.
- The concentration of the barium hydroxide solution is calculated from the mass of barium sulfate produced.

**Method 2**

- 25 cm<sup>3</sup> of the barium hydroxide solution is titrated with hydrochloric acid of known concentration.
- The concentration of the barium hydroxide solution is calculated from the result of the titration.

Compare the advantages and disadvantages of the two methods.

---



---



---



---



---



---



---



---



---



---




---

(5)

(Total 8 marks)

**Q18.**

(a) The diagram shows an atom of magnesium and an atom of chlorine.



Describe, in terms of electrons, how magnesium atoms and chlorine atoms change into ions to produce magnesium chloride ( $\text{MgCl}_2$ ).

---

---

---

---

---

---

---

---

---

---

(4)

(b) Calculate the relative formula mass ( $M_r$ ) of magnesium chloride ( $\text{MgCl}_2$ ).

Relative atomic masses ( $A_r$ ): magnesium = 24; chlorine = 35.5

---

---

---

Relative formula mass ( $M_r$ ) = \_\_\_\_\_

(2)

(Total 6 marks)


**Q19.**

A student investigated the rate of reaction of magnesium and hydrochloric acid.



The student studied the effect of changing the concentration of the hydrochloric acid.

She measured the time for the magnesium to stop reacting.



(a) The student changed the concentration of the hydrochloric acid.

Give **two** variables that the student should control.

1. \_\_\_\_\_
2. \_\_\_\_\_

(2)

(b) (i) The rate of reaction increased as the concentration of hydrochloric acid increased.

Explain why.

---

---

---

---

(2)

(ii) Explain why increasing the temperature would increase the rate of reaction.

---

---

---

---

---

---

(3)

- (c) (i) The student had a solution of sodium hydroxide with a concentration of 0.100 moles per dm<sup>3</sup>.

She wanted to check the concentration of a solution of hydrochloric acid.

She used a pipette to transfer 5.00 cm<sup>3</sup> of the hydrochloric acid into a conical flask.

She filled a burette with the 0.100 moles per dm<sup>3</sup> sodium hydroxide solution.

Describe how she should use titration to obtain accurate results.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

(4)

- (ii) Sodium hydroxide neutralises hydrochloric acid as shown in the equation:



The student found that 27.20 cm<sup>3</sup> of 0.100 moles per dm<sup>3</sup> sodium hydroxide neutralised 5.00 cm<sup>3</sup> of hydrochloric acid.

Calculate the concentration of the hydrochloric acid in moles per dm<sup>3</sup>.

Give your answer to three significant figures.

---

---

---

---

---

---

---

---

---

---

Concentration of hydrochloric acid = \_\_\_\_\_ moles per dm<sup>3</sup>

(3)



**Q20.**

Lead bromide is a solid. Some students were measuring how soluble lead bromide is at different temperatures.

This is the method they used.

- A** Pour 100 cm<sup>3</sup> of water into a beaker.
  - B** Heat or cool the water to the required temperature.
  - C** Add lead bromide to the water.
  - D** Stir until no more lead bromide dissolves.
  - E** Transfer 50 cm<sup>3</sup> of the lead bromide solution into an evaporating basin of known mass.
  - F** Heat the evaporating basin until all of the water has evaporated.
  - G** Measure the mass of the evaporating basin containing the dry lead bromide.
- (a) (i) How could the lead bromide solution be separated from the undissolved solid lead bromide after step **D**?

Draw a ring around the correct answer.

**electrolysis      filtration      neutralisation**

(1)

- (ii) Draw a ring around the correct answer to complete the sentence.

A suitable item of apparatus for measuring 50 cm<sup>3</sup> of the lead bromide solution

in step **E** is a measuring

cylinder.
funnel.
tube.

(1)

- (iii) One student's results are shown in **Table 1**.

**Table 1**

Volume of lead bromide solution	50 cm <sup>3</sup>
Mass of empty evaporating basin	35.4 g
Mass of the evaporating basin containing dry lead bromide	36.0 g

Calculate the mass of lead bromide dissolved in 50 cm<sup>3</sup> of lead bromide solution.

---

Mass of lead bromide dissolved = \_\_\_\_\_ g

(2)

(b) A different student got the results shown in **Table 2**.


**Table 2**

Temperature of lead bromide solution in °C	Mass of lead bromide dissolved in 50 cm <sup>3</sup> of solution in g
0	0.20
20	0.40
40	0.70
60	1.70
80	1.55
100	2.30

(i) Plot these results on the grid in **Graph 1**.

Draw a smooth curve of best fit.

**Graph 1**



(3)

(ii) One of the points is anomalous.

Draw a ring around the anomalous point on the graph.

Suggest **one** possible error in the experiment, and give a reason why this error would cause the anomalous point.

---

---

---

---


(3)

- (c) The solubility of lead bromide is so low that it can be made using a precipitation reaction.

A student investigated how much lead bromide was precipitated when different volumes of potassium bromide and lead nitrate solutions were mixed together.

This is the method the student used.

- Place  $10\text{ cm}^3$  of lead nitrate solution in a boiling tube.
- Using a burette, add  $2\text{ cm}^3$  of potassium bromide solution to the boiling tube containing the lead nitrate solution.
- Leave the mixture to stand.
- Measure the depth of the lead bromide precipitate using a ruler.
- Repeat using different volumes of potassium bromide solution.



- (i) A teacher suggested that the student should do the reaction in a measuring

cylinder.

Explain why it is a good idea to do the reaction in a measuring cylinder.


---

---

---

(2)

(ii) The student's results are plotted on **Graph 2**.



There are no anomalous points.

Complete the graph by drawing two straight lines through the points.

(2)

(iii) What depth of precipitate would you expect to get if 14 cm<sup>3</sup> of potassium bromide was used?

Give a reason for your answer.

Depth of precipitate \_\_\_\_\_ mm

Reason \_\_\_\_\_

---

---

(2)

- (iv) How would the results be different if the experiment was repeated using solutions at a higher temperature?

Give a reason for your answer.

---

---

---

---

---

(2)  
(Total 18 marks)


### Q21.

Some students were investigating the rate at which carbon dioxide gas is produced when metal carbonates react with an acid.

One student reacted 1.00 g of calcium carbonate with 50 cm<sup>3</sup>, an excess, of dilute hydrochloric acid.

The apparatus used is shown in **Diagram 1**.

**Diagram 1**



- (a) Complete the **two** labels for the apparatus on the diagram.
- (b) The student measured the volume of gas collected every 30 seconds.

(2)


The table shows the student's results.

Time in seconds	Volume of carbon dioxide collected in cm <sup>3</sup>
30	104
60	
90	198
120	221

150	232
180	238
210	240
240	240

- (i) **Diagram 2** shows what the student saw at 60 seconds.

**Diagram 2**



What is the volume of gas collected?

Volume of gas = \_\_\_\_\_ cm<sup>3</sup>

(1)

- (ii) Why did the volume of gas stop changing after 210 seconds?

---



---

(1)

- (c) Another student placed a conical flask containing 1.00 g of a Group 1 carbonate ( $M_2CO_3$ ) on a balance.

He then added 50 cm<sup>3</sup>, an excess, of dilute hydrochloric acid to the flask and measured the mass of carbon dioxide given off.

The equation for the reaction is:



The final mass of carbon dioxide given off was 0.32 g.

- (i) Calculate the amount, in moles, of carbon dioxide in 0.32 g carbon dioxide.

Relative atomic masses ( $A_r$ ): C = 12; O = 16

---



---

\_\_\_\_\_

Moles of carbon dioxide = \_\_\_\_\_ moles

(2)

- (ii) How many moles of the metal carbonate are needed to make this number of moles of carbon dioxide?

\_\_\_\_\_  
\_\_\_\_\_

Moles of metal carbonate = \_\_\_\_\_ moles

(1)

- (iii) The mass of metal carbonate used was 1.00 g.

Use this information, and your answer to part **(c) (ii)**, to calculate the relative formula mass ( $M_r$ ) of the metal carbonate.

If you could not answer part **(c) (ii)**, use 0.00943 as the number of moles of metal carbonate. This is **not** the answer to part **(c) (ii)**.

\_\_\_\_\_  
\_\_\_\_\_

Relative formula mass ( $M_r$ ) of metal carbonate = \_\_\_\_\_

(1)

- (iv) Use your answer to part **(c) (iii)** to calculate the relative atomic mass ( $A_r$ ) of the metal in the metal carbonate ( $M_2CO_3$ ) and so identify the Group 1 metal in the metal carbonate.

If you could not answer part **(c) (iii)**, use 230 as the relative formula mass of the metal carbonate. This is **not** the answer to part **(c) (iii)**.

To gain full marks, you must show your working.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Relative atomic mass of metal is \_\_\_\_\_

Identity of metal \_\_\_\_\_

(3)

- (d) Two other students repeated the experiment in part **(c)**.

- (i) When the first student did the experiment some acid sprayed out of the flask as the metal carbonate reacted.

Explain the effect this mistake would have on the calculated relative atomic mass of the metal.

---

---

---

---

---

---

---

---

(3)

- (ii) The second student used 100 cm<sup>3</sup> of dilute hydrochloric acid instead of 50 cm<sup>3</sup>.

Explain the effect, if any, this mistake would have on the calculated relative atomic mass of the metal.

---

---

---

---

---

---

---

---


(3)

(Total 17 marks)

**Q22.**

Magnesium reacts with steam to produce hydrogen gas and magnesium oxide.

A teacher demonstrated the reaction to a class. The figure below shows the apparatus the teacher used.



- (a) (i) The hydrogen produced was collected.

Describe how to test the gas to show that it is hydrogen.

Test \_\_\_\_\_



Result \_\_\_\_\_

\_\_\_\_\_ (2)

(ii) Explain why the magnesium has to be heated to start the reaction.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

(2)

(b) The equation for the reaction is:



(i) The teacher used 1.00 g of magnesium.

Use the equation to calculate the maximum mass of magnesium oxide produced.

Give your answer to three significant figures.

Relative atomic masses ( $A_r$ ): O = 16; Mg = 24

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Maximum mass = \_\_\_\_\_ g

(3)

(ii) The teacher's demonstration produced 1.50 g of magnesium oxide.

Use your answer from part (b)(i) to calculate the percentage yield.

If you could not answer part (b)(i), use 1.82 g as the maximum mass of magnesium oxide. This is **not** the answer to part (b)(i).

\_\_\_\_\_

Percentage yield = \_\_\_\_\_ %

(2)

(iii) Give **one** reason why the percentage yield is less than 100%.

\_\_\_\_\_  
\_\_\_\_\_

**Q23.**

Scientists found that a compound contained:

22.8% sodium; 21.8% boron; and 55.4% oxygen.

Use the percentages to calculate the empirical formula of the compound.

Relative atomic masses ( $A_r$ ): B = 11; O = 16; Na = 23

To gain full marks you **must** show all your working.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---


Empirical formula = \_\_\_\_\_

(Total 5 marks)

**Q24.**

A company manufactures ethanol ( $C_2H_5OH$ ).

The reaction for the process is:



The temperature and pressure can be changed to increase the yield of ethanol at equilibrium.

- (a) Explain what is meant by equilibrium.

---

---

---

---

---

---

(3)

- (b) (i) How would increasing the temperature change the **yield** of ethanol at equilibrium?

Give a reason for your answer.

---

---

---

---

---

(2)

- (ii) How would increasing the pressure change the **yield** of ethanol at equilibrium?

Give a reason for your answer.

---

---

---

---

---

(2)

- (c) A catalyst is added to increase the rate of the reaction.

Explain how adding a catalyst increases the rate of a chemical reaction.

---

---

---


---

(2)

(Total 9 marks)


**Q25.**

Fertilisers contain elements that plants need.



(a) **Figure 1** represents a nitrogen atom.

**Figure 1**



Complete each sentence.

(i) The mass number of this nitrogen atom is \_\_\_\_\_ .

(1)

(ii) Atoms of nitrogen with different numbers of neutrons are called

\_\_\_\_\_ .

(1)


(iii) Compared with a proton, the mass of an electron is

\_\_\_\_\_ .

(1)

(b) Fertilisers can be made from ammonia.

(i) Which diagram, **A**, **B**, or **C**, represents the electronic structure of an ammonia molecule?



(1)

The electronic structure of an ammonia molecule is shown in diagram

(ii) What is the correct formula of ammonia?

Draw a ring around the correct answer.



(1)

(c) A student made ammonium nitrate by reacting ammonia solution with an acid.

(i) Name the acid used to make ammonium nitrate.

\_\_\_\_\_

(1)

(ii) Complete the sentence.

The student added a few drops of \_\_\_\_\_, which changed colour

when the ammonia solution had neutralised the acid.

(1)

(iii) The student added charcoal and filtered the mixture.

This produced a colourless solution of ammonium nitrate.

How is solid ammonium nitrate obtained from the solution?

\_\_\_\_\_

(1)

(iv) A farmer put ammonium nitrate fertiliser onto a field of grass.

Suggest what would happen to the grass.

\_\_\_\_\_

\_\_\_\_\_

(1)

(d) Some fertilisers contain potassium chloride.


Potassium reacts with chlorine to produce potassium chloride.

**Figure 2** shows how this happens.

The dots (•) and crosses (x) represent electrons.

Only the outer shell is shown.

**Figure 2**



Use **Figure 2** to help you answer this question.

Describe, as fully as you can, what happens when potassium reacts with chlorine to produce potassium chloride.

---



---



---



---



---



---



---



---



---




---

(4)  
(Total 13 marks)

**Q26.**

Some students investigated reactions to produce magnesium.

- (a) The students used electrolysis to produce magnesium from magnesium chloride, as shown in the figure below.



- (i) Magnesium chloride contains magnesium ions and chloride ions.

Why does solid magnesium chloride **not** conduct electricity?

---



---

(1)

- (ii) One of the products of the electrolysis of molten magnesium chloride is magnesium.

Name the other product.

---

(1)

(iii) Why do magnesium ions ( $\text{Mg}^{2+}$ ) move to the negative electrode?

---

---

(1)

(iv) At the negative electrode, the magnesium ions ( $\text{Mg}^{2+}$ ) gain electrons to become magnesium atoms.

How many electrons does each magnesium ion gain?

---

(1)

(b) The students did the experiment four times and weighed the magnesium produced.

The table below shows their results.

Experiment	Mass of magnesium produced in grams
1	1.13
2	0.63
3	1.11
4	1.09

(i) There is an anomalous result.

Suggest **one** possible reason for the anomalous result.

---

---

(1)

(ii) Calculate the mean mass of magnesium produced, taking account of the anomalous result.

---

---

---

Mean mass = \_\_\_\_\_ g

(2)

(c) The formula of magnesium chloride is  $\text{MgCl}_2$

The relative formula mass of magnesium chloride is 95.

The relative atomic mass of magnesium is 24.

- (i) Use the equation to calculate the percentage mass of magnesium in magnesium chloride.

$$\text{Percentage mass of magnesium} = \frac{\text{mass of magnesium}}{\text{mass of magnesium chloride}} \times 100\%$$

---

---

---

Percentage mass of magnesium in magnesium chloride = \_\_\_\_\_ %

(2)


- (ii) Draw a ring around the relative mass of chlorine in  $\text{MgCl}_2$

71                      95                      119

(1)

- (d) Magnesium is also produced from the reaction of magnesium oxide with silicon.

- (i) The equation for the reaction is:



What is the meaning of this symbol  $\rightleftharpoons$  ?

Draw a ring around the correct answer.

**neutralisation reaction**

**precipitation reaction**

**reversible reaction**

(1)

- (ii) The forward reaction is endothermic.

Draw a ring around the correct answer to complete the sentence.

In an endothermic reaction the temperature of the surroundings

decreases.
increases.
stays the same.

(1)


(Total 12 marks)

### Q27.

This question is about zinc and magnesium.

Zinc is produced by electrolysis of molten zinc chloride, as shown in the figure below.





(a) (i) Why must the zinc chloride be molten for electrolysis?

---



---

(1)

(ii) Describe what happens at the negative electrode.

---



---



---



---



---

(3)

(iii) Complete the half equation for the reaction at the positive electrode.



(1)

(b) Magnesium can be produced from magnesium oxide.

The equation for the reaction is:



(i) How can you tell from the equation that the reaction is done at a high temperature?

---



---

(1)

(ii) This reaction to produce magnesium from magnesium oxide is **endothermic**.

What is meant by an **endothermic** reaction?

---



---

(1)

- (iii) A company made magnesium using this reaction.

Calculate the mass of magnesium oxide needed to produce 1.2 tonnes of magnesium.

Relative atomic masses ( $A_r$ ): O = 16; Mg = 24

---

---

---

---

Mass of magnesium oxide needed = \_\_\_\_\_ tonnes

(3)

- (iv) The company calculated that they would produce 1.2 tonnes of magnesium, but only 0.9 tonnes was produced.

Calculate the percentage yield.

---

---

Percentage yield = \_\_\_\_\_ %

(1)

- (v) Give **one** reason why the calculated yield of magnesium might not be obtained.

---

---

(1)

(Total 12 marks)

### Q28.

Thermosoftening polymers can be used to make plastic bottles and food packaging.

- (a) Why are thermosoftening polymers **not** suitable for storing very hot food?

---

---

(1)

- (b) The reaction to produce the polymers uses a catalyst.

Why are catalysts used in chemical reactions?

---

---

(1)

- (c) Compounds from food packaging must not get into food.

Gas chromatography can be used to separate compounds in food.

The output from the gas chromatography column can be linked to an instrument which can identify the compounds.

- (i) Name the instrument used to identify the compounds.

---

---

(1)

- (ii) Give **one** reason why instrumental methods of analysis are used to identify the compounds.

---

---

(1)

- (d) Poly(ethene) is a thermosoftening polymer.

Poly(ethene) can be made with different properties. The properties depend on the conditions used when poly(ethene) is made.

Suggest **two** conditions which could be changed when poly(ethene) is made.

---

---

(2)

(Total 6 marks)

**Q29.**

Printed pictures can be made using etchings.



© Eduardo Jose Bernardino/iStock

An etching can be made when a sheet of brass reacts with iron chloride solution.

(a) Brass is a mixture of two metals, copper and zinc.

(i) A mixture of two metals is called \_\_\_\_\_ .

(1)

(ii) Draw a ring around the correct answer to complete the sentence.

Copper and zinc atoms are different sizes.

This makes brass 

harder
more flexible
softer

 than the pure metals.

(1)

(b) Iron chloride has the formula  $\text{FeCl}_3$

Relative atomic masses ( $A_r$ ): Cl = 35.5; Fe = 56.

(i) Calculate the relative formula mass ( $M_r$ ) of iron chloride ( $\text{FeCl}_3$ ).

---



---



---

Relative formula mass ( $M_r$ ) of iron chloride = \_\_\_\_\_

(2)

(ii) Calculate the percentage of iron in iron chloride ( $\text{FeCl}_3$ ).

Percentage of iron in iron chloride = \_\_\_\_\_%

(2)


(Total 6 marks)

**Q30.**

This question is about compounds of copper.

- (a) A student made some copper(II) sulfate crystals.

The flow diagram shows the stages of the preparation of copper(II) sulfate crystals.




- (i) The reaction mixture is heated in **Stage 1**.

Suggest why.

\_\_\_\_\_  
\_\_\_\_\_

(1)

- (ii) Complete the equation for this reaction.



(2)

- (iii) How would the student remove the unreacted copper(II) oxide in **Stage 2**?

\_\_\_\_\_  
\_\_\_\_\_

(1)

- (iv) How would the student obtain copper(II) sulfate crystals from the copper(II) sulfate solution in **Stage 3**?

\_\_\_\_\_

(1)

- (v) The mass of crystals obtained was less than the student had calculated.


Suggest **one** reason why.

\_\_\_\_\_  
\_\_\_\_\_

(1)

(b) The student heated the blue copper(II) sulfate crystals.

The word equation for the reaction is shown below.



(i) What does the symbol  $\rightleftharpoons$  mean ?

\_\_\_\_\_

(1)

(ii) 300 J of energy are taken in when some blue copper(II) sulfate crystals are heated.

What is the energy change when an excess of water is added to the anhydrous copper(II) sulfate produced?

\_\_\_\_\_

\_\_\_\_\_

(2)

(c) A sample of copper nitride contains 3.81 g of copper and 0.28 g of nitrogen.

Calculate the empirical formula.

You **must** show all your working to get full marks.

Relative atomic masses ( $A_r$ ): N = 14; Cu = 63.5.

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_


Empirical formula = \_\_\_\_\_

(4)

(Total 13 marks)

### Q31.

Etching is a way of making printed circuit boards for computers.



© Dario Lo Presti/Shutterstock

Printed circuit boards are made when copper sheets are etched using iron(III) chloride solution. Where the copper has been etched, only plastic remains.

- (a) Copper is a good conductor of electricity.

Explain why.

---


---

---

---

(2)

- (b) Iron(III) chloride can be produced by the reaction shown in the equation:



- (i) Calculate the maximum mass of iron(III) chloride ( $\text{FeCl}_3$ ) that can be produced from 11.20 g of iron.

Relative atomic masses ( $A_r$ ): Cl = 35.5; Fe = 56.

---

---

---

---

---

Maximum mass of iron(III) chloride = \_\_\_\_\_ g

(3)

- (ii) The actual mass of iron(III) chloride ( $\text{FeCl}_3$ ) produced was 24.3 g.

Calculate the percentage yield.

(If you did not answer part (b)(i) assume that the maximum theoretical mass of iron(III) chloride ( $\text{FeCl}_3$ ) is 28.0 g. This is **not** the correct answer to part (b)(i).)


Percentage yield = \_\_\_\_\_ %

(1)


(Total 6 marks)

**Q32.**

This apparatus is used for the reaction of copper oxide (CuO) with methane (CH<sub>4</sub>).



- (a) The symbol equation for this reaction is shown below.



The water and carbon dioxide produced escape from the test tube.

Use information from the equation to explain why.

---

---

(1)

- (b) (i) Calculate the relative formula mass ( $M_r$ ) of copper oxide (CuO).

Relative atomic masses ( $A_r$ ): O = 16, Cu = 64

---

---

---

Relative formula mass ( $M_r$ ) = \_\_\_\_\_

(2)

- (ii) Calculate the percentage of copper in copper oxide.

---

---

---

Percentage of copper = \_\_\_\_\_ %

(2)

- (iii) Calculate the maximum mass of copper that could be produced from 4.0 g of



copper oxide.

---

---

Mass of copper produced = \_\_\_\_\_ g

(1)

- (c) The experiment was done three times.

The mass of copper oxide used and the mass of copper produced were measured each time.

The results are shown in the table.

	Experiment		
	1	2	3
Mass of copper oxide used in g	4.0	4.0	4.0
Mass of copper produced in g	3.3	3.5	3.2

- (i) Calculate the mean mass of copper produced in these experiments.

---

---

Mean mass of copper produced = \_\_\_\_\_ g

(1)

- (ii) Suggest how the results of the experiment could be made more precise.

---

---

(1)

- (iii) The three experiments gave different results for the amount of copper produced.

This was caused by experimental error.

Suggest two causes of experimental error in these experiments.

1. \_\_\_\_\_  
\_\_\_\_\_  
2. \_\_\_\_\_  
\_\_\_\_\_


(2)

(Total 10 marks)

**Q33.**

Ammonium sulfate and urea are made from ammonia. These compounds are used by farmers.

The flow diagram shows the stages to make ammonium sulfate and urea.



- (a) Give **two** examples from the flow diagram of the efficient use of energy and raw materials.

---



---



---



---




---



---

(2)

- (b) The equation for the reaction in Stage 4 is shown below.



The forward reaction is exothermic.

State **and** explain:

- (i) how a **decrease** in temperature would affect the yield of ammonia at equilibrium

---

---

---

---

---

(2)

- (ii) how an **increase** in pressure would affect the yield of ammonia at equilibrium.

---

---

---

---

---

(2)

- (c) The equation for the reaction in Stage 7 is shown below.



The table gives the relative formula masses ( $M_r$ ) of the reactants and the products for this reaction.

Formula of reactant or product	Relative formula masses ( $M_r$ )
NH <sub>3</sub>	17
CO <sub>2</sub>	44
NH <sub>2</sub> CONH <sub>2</sub>	60
H <sub>2</sub> O	18

Percentage atom economy can be calculated using:

$$\text{Percentage atom economy} = \frac{M_r \text{ of useful product}}{\text{total } M_r \text{ of all reactants added together}} \times 100\%$$

Calculate the percentage atom economy for the reaction in Stage 7.

---

---

---

---

---

---

Percentage atom economy = \_\_\_\_\_ %


(2)

(Total 8 marks)

**Q34.**

Saturated hydrocarbons, for example methane and octane, are often used as fuels.

(a) Methane can be represented as:



(i) The formula of methane is \_\_\_\_\_ .

(1)

(ii) Draw a ring around the correct answer to complete the sentence.

In a saturated hydrocarbon molecule all of the bonds are

double.
ionic.
single.

(1)

(iii) Draw a ring around the correct answer to complete the sentence.

The homologous series that contains methane and octane is called the

alcohols.
alkanes.
alkenes.

(1)

(b) (i) The complete combustion of petrol produces carbon dioxide, water vapour and sulfur dioxide.

Name **three** elements petrol must contain.

1. \_\_\_\_\_

2. \_\_\_\_\_

3. \_\_\_\_\_

(3)

(ii) The exhaust gases from cars can contain oxides of nitrogen.

Complete the sentence.

Nitrogen in the oxides of nitrogen comes from \_\_\_\_\_ .

(1)

(iii) The sulfur dioxide and oxides of nitrogen from cars cause an environmental problem.

Name the problem and describe **one** effect of the problem.

Name of problem \_\_\_\_\_

Effect of problem \_\_\_\_\_

\_\_\_\_\_

(2)

(c) When a fuel burns without enough oxygen, there is incomplete combustion.

One gaseous product of incomplete combustion is carbon monoxide.

Name **one** solid product of incomplete combustion.

\_\_\_\_\_


(1)

(d) A student investigated how well different hydrocarbon fuels would heat up 100 g of water.

Her hypothesis was:

**The more carbon atoms there are in a molecule of any fuel, the better the fuel is.**

The apparatus the student used is shown in the diagram.



She burned each hydrocarbon fuel for 2 minutes.

Her results are shown in the table.

Name of hydrocarbon fuel	Number of carbon atoms in a molecule of hydrocarbon	Temperature change of water in °C after 2	Temperature change per g of fuel burned	Observations
--------------------------	---	---	---	--------------

	<b>fuel</b>	<b>minutes</b>		
Pentane	5	30	60	no smoke
Hexane	6	40	57	very small amount of smoke
Octane	8	55	55	small amount of smoke
Decane	10	57	52	large amount of smoke
Dodecane	12	60	43	very large amount of smoke

The student investigated only hydrocarbons.

Look carefully at her results.

How well do the student's results support her hypothesis?

**The more carbon atoms there are in a molecule of any fuel, the better the fuel is.**

Give reasons for your answer.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

(4)

(e) A 0.050 mol sample of a hydrocarbon was burned in excess oxygen.

The products were 3.60 g of water and 6.60 g of carbon dioxide.

(i) Calculate the number of moles of carbon dioxide produced.

Relative atomic masses: C = 12; O = 16.

---

---

Moles of carbon dioxide = \_\_\_\_\_

(2)

- (ii) When the hydrocarbon was burned 0.20 mol of water were produced.

How many moles of hydrogen atoms are there in 0.20 mol of water?

\_\_\_\_\_

Moles of hydrogen atoms = \_\_\_\_\_

(1)

- (iii) The amount of hydrocarbon burned was 0.050 mol.

Use this information and your answers to parts **(e) (i)** and **(e) (ii)** to calculate the molecular formula of the hydrocarbon.

If you could not answer parts **(e) (i)** or **(e) (ii)** use the values of 0.20 moles carbon dioxide and 0.50 moles hydrogen. These are **not** the answers to parts **(e) (i)** and **(e) (ii)**.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Formula = \_\_\_\_\_

(2)

(Total 19 marks)

**Q35.**

- (a) A student had a colourless solution.

The student thought the solution was dilute hydrochloric acid.

- (i) The student added universal indicator to this solution.

What colour would the universal indicator change to if the solution is hydrochloric acid?

\_\_\_\_\_


(1)

- (ii) Describe how the student could show that there are chloride ions in this solution.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

(2)

(b) The results of a titration can be used to find the concentration of an acid.



Describe how to use the apparatus to do a titration using 25 cm<sup>3</sup> of dilute hydrochloric acid.

In your answer you should include:

- how you will determine the end point of the titration
- how you will make sure the result obtained is accurate.

---

---

---

---

---

---

---

---

---

---

---

---

---

**(4)**

(c) Hydrochloric acid is a strong acid.

Ethanoic acid is a *weak acid*.

What is meant by the term *weak acid*?


---



---

(1)

(d) The displayed formula of ethanoic acid is:



(i) On the formula, draw a circle around the functional group in ethanoic acid.

(1)

(ii) Ethanoic acid and ethanol react together to make the ester ethyl ethanoate.

Draw the **displayed** formula of ethyl ethanoate.

(2)

(Total 11 marks)

### Q36.

This question is about lithium and sodium.

(a) Use the Chemistry Data Sheet to help you to answer this question.


In which group of the periodic table are lithium and sodium?

Group

(1)

(b) A lithium atom can be represented as  ${}^7_3\text{Li}$

The diagram represents the lithium atom.



- (i) Some particles in the nucleus have a positive charge.

What is the name of these particles?

\_\_\_\_\_

(1)

- (ii) Some particles in the nucleus have no charge.

What is the name of these particles?

\_\_\_\_\_

(1)

- (iii) Use the correct answer from the box to complete the sentence.

3	4	7
---	---	---

The mass number of this atom of lithium is


(1)

- (c) Sodium reacts with chlorine to produce sodium chloride.



The diagram shows how the reaction happens.

Only the outer electrons are shown.



Draw a ring around the correct answer to complete each sentence.

- (i) A sodium atom changes into a sodium ion by

gaining

losing

sharing

an electron.

(1)

(ii) A sodium ion has

a negative
no
a positive

charge.

(1)

(iii) The ions in sodium chloride are held together by

strong

covalent
electrostatic
magnetic

forces.

(1)

(d) Sodium chloride is an ionic compound.

Tick (✓) **two** properties of ionic compounds.

Property	Tick (✓)
Do <b>not</b> dissolve in water	
High melting points	
Low boiling points	
Strong bonds	

(2)

(e) (i) The formula of sodium chloride is NaCl

Calculate the relative formula mass of sodium chloride.

Relative atomic masses: Na = 23; Cl = 35.5

\_\_\_\_\_

\_\_\_\_\_

Relative formula mass = \_\_\_\_\_

(1)

(ii) Draw a ring around the correct answer to complete each sentence.

The relative formula mass of a substance, in grams,

is one

ion
isotope
mole

of the substance.

- (f) Nanoparticles of sodium chloride (salt) are used to flavour crisps.

What are nanoparticles?

---




---

(1)

(Total 12 marks)

**Q37.**

- (a) Some students did an experiment to find the temperature change when hydrochloric acid reacts with sodium hydrogencarbonate.



The results are in the table.

Number of spatula measures of sodium hydrogencarbonate	Start temperature in °C	Final temperature in °C	Change in temperature in °C
2	20	16	4
4	20	14	6
6	19	11	8
8	20	10	10
10	19	9	10
12	20	10	10

- (i) Describe, as fully as you can, the trends shown in the students' results.

---



---



---



---

---

---

---

---

(3)


(ii) State the type of energy transfer for this reaction.

---

---

(1)

- (b) Sodium hydrogencarbonate is used as baking powder for making cakes.  
When the cake mixture is baked the sodium hydrogencarbonate decomposes.  
The equation for the reaction is:



(i) The cake mixture rises when baked.



© Michael Valdez/iStock


Use the equation to suggest why.

---

---

(1)

(ii) The same reaction can be reversed to produce sodium hydrogencarbonate from sodium carbonate.



Do the reactants need to be heated?

Give a reason for your answer.

---

---

(1)

(c) (i) Calculate the relative formula mass of sodium hydrogencarbonate ( $\text{NaHCO}_3$ ).

Relative atomic masses ( $A_r$ ): H=1; C=12; O=16; Na=23

---

---

---

Relative formula mass ( $M_r$ ) = \_\_\_\_\_

(2)

(ii) Calculate the percentage by mass of carbon in sodium hydrogencarbonate.

---

---

Percentage of carbon = \_\_\_\_\_ %


(1)

(Total 9 marks)

**Q38.**

Ammonia is produced from nitrogen and hydrogen.

The equation for this reaction is:



(a) (i) A company wants to make 6.8 tonnes of ammonia.

Calculate the mass of nitrogen needed.

Relative atomic masses ( $A_r$ ): H = 1; N = 14

---

---

---

---

---

Mass of nitrogen = \_\_\_\_\_ tonnes

(3)

(ii) The company expected to make 6.8 tonnes of ammonia.

The yield of ammonia was only 4.2 tonnes.

Calculate the percentage yield of ammonia.

---

---

Percentage yield of ammonia = \_\_\_\_\_ %

(2)

- (iii) Use the equation above to explain why the percentage yield of ammonia was less than expected.


---

---

(1)

- (b) Complete the diagram to show the arrangement of the outer shell electrons of the nitrogen and hydrogen atoms in ammonia.

Use dots (•) and crosses (x) to represent the electrons.



(2)

- (c) Ammonia dissolves in water to produce an alkaline solution.

- (i) Which ion makes ammonia solution alkaline?

---

(1)

- (ii) Name the type of reaction between aqueous ammonia solution and an acid.

---

(1)

- (iii) Name the acid needed to produce ammonium nitrate.

---

(1)

- (iv) The reaction of ammonia with sulfuric acid produces ammonium sulfate.

Use the formulae of the ions on the Chemistry Data Sheet.

Write the formula of ammonium sulfate.

---


(1)

(Total 12 marks)

**Q39.**

In this question you will be assessed on using good English, organising information clearly and using specialist terms where appropriate.

A student used the equipment shown to do a titration.



Describe how the student should use this equipment to find the volume of sodium hydroxide solution that reacts with a known volume of acid. Include any measurements the student should make.

Do **not** describe how to do any calculations.

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---



---

---

---

(Total 6 marks)

**Q40.**

Some students investigated magnesium oxide.

(a) Magnesium oxide has the formula MgO.

(i) Calculate the relative formula mass ( $M_r$ ) of magnesium oxide.

Relative atomic masses: O = 16; Mg = 24.

---

---

Relative formula mass = \_\_\_\_\_

(2)

(ii) Calculate the percentage by mass of magnesium in magnesium oxide.

---

---

Percentage by mass of magnesium in magnesium oxide = \_\_\_\_\_%

(2)

(iii) Calculate the mass of magnesium needed to make 25 g of magnesium oxide.


---

Mass of magnesium = \_\_\_\_\_ g

(1)

(b) The students calculated that if they used 0.12 g of magnesium they should make 0.20 g of magnesium oxide.

They did this experiment to find out if this was correct.



- The students weighed 0.12 g of magnesium ribbon into a crucible.
- They heated the magnesium ribbon.
- They lifted the lid of the crucible slightly from time to time to allow air into the crucible.
- The students tried to avoid lifting the lid too much in case some of the magnesium oxide escaped.
- When all of the magnesium appeared to have reacted, the students weighed the magnesium oxide produced.

The results of the experiment are shown below.

Mass of magnesium used in grams	0.12
Mass of magnesium oxide produced in grams	0.18

- (i) The mass of magnesium oxide produced was lower than the students had calculated.  
They thought that this was caused by experimental error.

Suggest **two** experimental errors that the students had made.

---



---



---



---

(2)

- (ii) The students only did the experiment once.

Give **two** reasons why they should have repeated the experiment.

---

---


---

---

(2)  
(Total 9 marks)

**Q41.**

The symbol equation for the decomposition of hydrogen peroxide is:



- (a) This reaction is *exothermic*.


What is an *exothermic* reaction?

---


---

(1)

- (b) A student measured the volume of oxygen produced by 50 cm<sup>3</sup> of hydrogen peroxide.



The graph shows the results.



- (i) Use the graph to describe the changes in the rate of the reaction from 0 to 35 seconds.

---

---

---

---

---

---

---

(3)

- (ii) What was the total volume of oxygen gas collected?

\_\_\_\_\_ cm<sup>3</sup>

(1)

- (iii) The student had calculated that the hydrogen peroxide used should produce 25 cm<sup>3</sup> of oxygen.

Calculate the percentage yield of oxygen.

---

---

---

Answer = \_\_\_\_\_ %

(2)

- (c) An increase in the temperature of the hydrogen peroxide increases the rate of the reaction.

Use your knowledge of particles to explain why.

---

---

---

---

---

---

---

(3)

(Total 10 marks)

### Q42.

Aluminium is extracted from aluminium oxide.

- (a) The formula of aluminium oxide is Al<sub>2</sub>O<sub>3</sub>

The relative formula mass ( $M_r$ ) of aluminium oxide is 102.

Calculate the percentage of aluminium in aluminium oxide.

Relative atomic masses ( $A_r$ ): O = 16; Al = 27.

---

---

---


---

Percentage of aluminium = \_\_\_\_\_ %

(2)

- (b) Aluminium is extracted from aluminium oxide using electrolysis.

The diagram shows a cell used for the extraction of aluminium.



- (i) The electrolyte contains cryolite.

Explain why.

---


---

---

---

(2)

- (ii) Oxygen is formed at the positive electrode. Complete and balance the equation for this reaction.



(2)

- (iii) The positive electrode in the cell is used up during the process.

Explain why.

---

---

---

---

---

---

(2)  
(Total 8 marks)

**Q43.**

This question is about calcium hydroxide.

Ancient artworks and monuments can be protected from acid rain if the surface is sprayed with calcium hydroxide nanoparticles.



By Svilen Enev (Own work) [GFDL or CC-BY-SA-3.0], via Wikimedia Commons

- (a) Calcium hydroxide has the formula  $\text{Ca}(\text{OH})_2$

Why are there two hydroxide ions for each calcium ion in the formula?

---

---

---

(1)

- (b) The calcium hydroxide is used in the form of *nanoparticles*.

What are *nanoparticles*?

---

---

(1)

- (c) A student added water to calcium oxide to make calcium hydroxide.

The equation for the reaction is shown below.



Calculate the maximum mass of calcium hydroxide which could be made from 2.00 g of calcium oxide.

Relative atomic masses ( $A_r$ ): H = 1; O = 16; Ca = 40.

---

---

---

---

---

---

---


Maximum mass of calcium hydroxide = \_\_\_\_\_ g

(3)

(Total 5 marks)

#### Q44.

- (a) Calcium chloride is made from limestone. Limestone contains mainly calcium carbonate and a small amount of magnesium carbonate.



- (i) In **stage 1** calcium carbonate reacts with acid **X** to form calcium chloride.

Draw a ring around the name of acid **X**.

hydrochloric

nitric

sulfuric

(1)

- (ii) **Stage 1** produces a concentrated solution of calcium chloride. The solution also contains magnesium chloride.

Calcium hydroxide solution is added in **stage 2** to remove the magnesium chloride.

The equation for this reaction is:



Draw a ring around the correct answer to complete each sentence.

In **stage 2** a precipitate is made because

magnesium hydroxide is 

dissolved
insoluble
soluble

 in water.

In **stage 3** the solid magnesium hydroxide can be separated from the calcium

chloride solution using 

chromatography.
electrolysis.
filtration.

(2)

- (iii) What method can be used to change the calcium chloride solution into solid calcium chloride?  
Draw a ring around your answer.

**crystallisation**

**electrolysis**

**reduction**

(1)


- (b) Calcium chloride can also be made by reacting calcium with chlorine:



The diagram shows what happens to atoms of calcium and chlorine in this reaction.

The dots (•) and crosses (x) are used to represent electrons.

Only the outer electrons are shown.



Use the diagram to help you to answer this question.



Describe, as fully as you can, what happens when calcium reacts with chlorine to make calcium chloride.

---

---

---

---

---

---

---

---


---

---

(4)  
(Total 8 marks)

**Q45.**

Some students did an experiment to find the relative formula mass ( $M_r$ ) of a gas.



This is the method they used.

- The mass of the canister of gas was measured using a balance, which weighed to two decimal places.
- The measuring cylinder was filled with 1 dm<sup>3</sup> of the gas from the canister.
- The mass of the canister of gas was measured again.
- The temperature of the laboratory was measured.
- The air pressure in the laboratory was measured.

The students repeated the experiment three times.

(a) The results for one of the experiments are shown in the table below.

Mass of the canister of gas before filling the measuring cylinder	53.07 g
---	---------

Mass of the canister of gas after filling the measuring cylinder	51.21 g
--	---------

Calculate the mass of the 1 dm<sup>3</sup> of gas in the measuring cylinder.

Mass = \_\_\_\_\_ g

(1)

- (b) How could the results be made more precise?

---



---

(1)

- (c) The students used their results to calculate values for the relative formula mass ( $M_r$ ) of this gas.  
The results are shown in the table below.

Experiment	1	2	3	4
Relative formula mass ( $M_r$ )	45.4	51.5	46.3	45.8

- (i) Calculate the mean value for these results.

Mean = \_\_\_\_\_

(2)

- (ii) The four results are different.  
The students thought this was because of experimental error.

Suggest **two** causes of experimental error in this experiment.

---



---



---



---

(2)

- (iii) It was important for the students to repeat the experiment.  
Suggest why.

---



---

(1)

- (d) The teacher told the students that the formula of the gas is C<sub>3</sub>H<sub>8</sub>

Calculate the relative formula mass ( $M_r$ ) of this gas. You should show your working.

Relative atomic masses: H = 1; C = 12.

---


---

Relative formula mass = \_\_\_\_\_

(2)  
(Total 9 marks)

**Q46.**

Ammonia is made using the Haber process.



- (a) How is ammonia separated from unreacted nitrogen and hydrogen in the separator?

---


---

---

---

(2)

- (b) The equation shows the reaction which takes place in the reactor:



- (i) Why does the yield of ammonia at equilibrium increase as the temperature is decreased?

---

---

(1)

- (ii) A temperature of 450 °C is used in the reactor to make the reaction take place quickly.

Explain, in terms of particles, why increasing the temperature makes a reaction go faster.

---

---

---

---

(2)

- (iii) Why does the yield of ammonia at equilibrium increase as the pressure is increased?

---

---

(1)


- (iv) The pressure used in the reactor is 200 atmospheres. Suggest why a much higher pressure is **not** used.

---

---

(1)

- (c) Use the equation for the reaction in the reactor to help you to answer these questions.



- (i) It is important to mix the correct amounts of hydrogen and nitrogen in the reactor.

20 m<sup>3</sup> of nitrogen is reacted with hydrogen.

What volume of hydrogen (measured at the same temperature and pressure as the nitrogen) is needed to have the correct number of molecules to react with the nitrogen?

Volume of hydrogen needed = \_\_\_\_\_ m<sup>3</sup>

(1)

- (ii) Calculate the maximum mass of ammonia that can be made from 2 g of nitrogen.

Relative atomic masses: H = 1; N = 14.

---

---

---

---

---

---

Maximum mass of ammonia = \_\_\_\_\_ g

(3)

(d) The expected maximum mass of ammonia produced by the Haber process can be calculated.

(i) In one process, the maximum mass of ammonia should be 80 kg.

The actual mass of ammonia obtained was 12 kg.

Calculate the percentage yield of ammonia in this process.

---

---

Percentage yield of ammonia = \_\_\_\_\_ %

(1)

(ii) Give **two** reasons why it does **not** matter that the percentage yield of ammonia is low.

Use the flow diagram at the start of this question to help you.

---

---

---

---

(2)

(Total 14 marks)

### Q47.

Vinegar can be added to food.

Vinegar is a solution of ethanoic acid in water.



(a) Ethanoic acid is a *weak* acid.

Draw a ring around the correct answer to complete each sentence.

(i) When dissolved in water, an acid forms a solution containing

carbonate ions.

hydrogen ions.

hydroxide ions.

(1)

(ii) Ethanoic acid is a *weak* acid because in water it is


completely ionised.

not ionised.

partially ionised.

(1)

(b) The diagram shows the apparatus used to investigate the amount of ethanoic acid in vinegar.



(i) Draw a ring around the name of the piece of apparatus labelled **A** on the diagram.

burette

measuring cylinder

pipette

(1)

(ii) Phenolphthalein is added to the vinegar in the conical flask so that the end point of the titration can be seen.

What type of substance is phenolphthalein?

Draw a ring around the correct answer.

**alkali**                      **catalyst**                      **indicator**

(1)

(iii) How would you know that the end point of the titration has been reached?

---

---

(1)

(c) The results of the titration are shown in the table.

	Rough titration	Accurate titrations		
		1	2	3
Final reading in cm <sup>3</sup>	22	21.30	22.50	24.40
Initial reading in cm <sup>3</sup>	0	1.00	2.00	4.00
Volume used in cm <sup>3</sup>	22	20.30	20.50	20.40

Calculate the best value of the mean volume from these titrations.

---

---

---

Mean volume used = \_\_\_\_\_ cm<sup>3</sup>

(2)

(d) 25.0 cm<sup>3</sup> of this vinegar contained 1.25 g of ethanoic acid.

Calculate the mass of ethanoic acid in 1 litre (1000 cm<sup>3</sup>) of this vinegar.

---

---

---

Mass = \_\_\_\_\_ g

(2)

(Total 9 marks)

**Q48.**

Vinegar can be added to food. Vinegar is an aqueous solution of ethanoic acid.



Ethanoic acid is a *weak* acid.

- (a) Which ion is present in aqueous solutions of all acids?

---

(1)

- (b) What is the difference between the pH of a *weak* acid compared to the pH of a strong acid of the same concentration?

Give a reason for your answer.

---


---

---

---

(2)

- (c) The diagram shows the apparatus used to find the concentration of ethanoic acid in vinegar.



- (i) Why should phenolphthalein indicator be used for this titration instead of methyl orange?



---

---

(1)

- (ii) 25.00 cm<sup>3</sup> of vinegar was neutralised by 30.50 cm<sup>3</sup> of a solution of sodium hydroxide with a concentration of 0.50 moles per cubic decimetre.

The equation for this reaction is:



Calculate the concentration of ethanoic acid in this vinegar.

---

---

---

---

Concentration of ethanoic acid in this vinegar = \_\_\_\_\_ moles per cubic decimetre

(2)

- (d) The concentration of ethanoic acid in a different bottle of vinegar was 0.80 moles per cubic decimetre.

Calculate the mass in grams of ethanoic acid (CH<sub>3</sub>COOH) in 250 cm<sup>3</sup> of this vinegar. The relative formula mass (*M<sub>r</sub>*) of ethanoic acid = 60.

---

---

---

---

Mass of ethanoic acid = \_\_\_\_\_ g

(2)

(Total 8 marks)

**Q49.**

Calamine lotion is used to treat itching. The main ingredients are two metal oxides.



- (a) One of the metal oxides has a relative formula mass ( $M_r$ ) of 81.

The formula of this metal oxide is MO.  
(M is **not** the correct symbol for the metal.)

The relative atomic mass ( $A_r$ ) of oxygen is 16.

- (i) Calculate the relative atomic mass ( $A_r$ ) of metal M.

---

---

---

Relative atomic mass ( $A_r$ ) = \_\_\_\_\_

(2)

- (ii) Use your answer to part (a)(i) and the periodic table on the Data Sheet to name metal M.

The name of metal M is \_\_\_\_\_ .

(1)

- (b) The other metal oxide is iron(III) oxide.

This contains iron(III) ions ( $\text{Fe}^{3+}$ ) and oxide ions ( $\text{O}^{2-}$ ).

- (i) Explain in terms of electrons how an iron atom (Fe) can change into an iron(III) ion ( $\text{Fe}^{3+}$ ).

---


---

---


---

(2)

- (ii) The diagram below represents the electronic structure of an oxygen atom (O).




Complete the diagram below to show the electronic structure of an oxide ion ( $O^{2-}$ ).




(1)  
(Total 6 marks)

**Q50.**

An experiment was done on the reaction of copper oxide ( $CuO$ ) with methane ( $CH_4$ ).



(a) The equation for this reaction is shown below.



The water and carbon dioxide produced escapes from the test tube.

Use information from the equation to explain why.

(1)

(b) (i) Calculate the relative formula mass ( $M_r$ ) of copper oxide ( $CuO$ ).

Relative atomic masses ( $A_r$ ): O = 16; Cu = 64.

---



---

Relative formula mass ( $M_r$ ) = \_\_\_\_\_

(2)

- (ii) Calculate the percentage of copper in copper oxide.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

Percentage of copper = \_\_\_\_\_ %

(2)

- (iii) Calculate the mass of copper that could be made from 4.0 g of copper oxide.

\_\_\_\_\_  
\_\_\_\_\_

Mass of copper = \_\_\_\_\_ g

(1)

- (c) The experiment was done three times.  
The mass of copper oxide used and the mass of copper made was measured each time.  
The results are shown in the table.

	Experiment		
	1	2	3
Mass of copper oxide used in g	4.0	4.0	4.0
Mass of copper made in g	3.3	3.5	3.2

- (i) Calculate the mean mass of copper made in these experiments.

\_\_\_\_\_  
\_\_\_\_\_

Mean mass of copper made = \_\_\_\_\_ g

(1)

- (ii) Suggest how the results of these experiments could be made more precise.

\_\_\_\_\_  
\_\_\_\_\_

(1)

- (iii) The three experiments gave slightly different results for the mass of copper made.  
This was caused by experimental error.

Suggest **two** causes of experimental error in these experiments.

1. \_\_\_\_\_  
\_\_\_\_\_
2. \_\_\_\_\_  
\_\_\_\_\_

(2)  
(Total 10 marks)

### Q51.

Read the information about protecting the bottoms of ships.

#### A Copper-bottomed Investment



From the 16th to the 19th century, the bottoms of many wooden ships were protected from marine organisms by being covered with sheets of metal.

At first lead was used on the bottoms of ships, then copper was used until 1832 when Muntz Metal replaced it. Muntz Metal is an alloy of two transition metals, copper and zinc.

**Table of data**

	<b>Lead</b>	<b>Copper</b>	<b>Muntz Metal</b>
<b>Cost (£/kg)</b>	£1.20	£3.20	£2.30
<b>Melting point (°C)</b>	327	1083	904
<b>Stops sea worms attacking wood</b>	Yes	Yes	Yes
<b>Stops barnacles and seaweed sticking to the bottom of the ship</b>	No	Yes	Yes

(a) Use the information to answer the following questions.

(i) Suggest why copper replaced lead.

---

---

(1)

(ii) Suggest why Muntz Metal replaced copper.

---

---

(1)

(b) A sample of Muntz Metal contains a very small amount of iron as an impurity.

(i) Name an instrumental method of analysis that could be used to detect iron.

---

(1)

(ii) Suggest why an instrumental method would detect the iron in this sample of Muntz Metal but a chemical method is **not** likely to be successful.

---

---

(1)

(c) Today, ships are made from steel. Steels are alloys of iron, a transition metal.

Give **two** properties of transition metals that make them suitable for making ships.

Property 1 \_\_\_\_\_

---

Property 2 \_\_\_\_\_


---

(2)

(Total 6 marks)

**Q52.**

Ammonium sulfate is an artificial fertiliser.



- (a) (i) When this fertiliser is warmed with sodium hydroxide solution, ammonia gas is given off.  
Describe and give the result of a test for ammonia gas.

Test \_\_\_\_\_

\_\_\_\_\_

Result \_\_\_\_\_

\_\_\_\_\_

(2)

- (ii) Describe and give the result of a chemical test to show that this fertiliser contains sulfate ions ( $\text{SO}_4^{2-}$ ).

Test \_\_\_\_\_

\_\_\_\_\_

Result \_\_\_\_\_

\_\_\_\_\_

(2)

- (b) Ammonium sulfate is made by reacting sulfuric acid (a *strong* acid) with ammonia solution (a *weak* alkali).

- (i) Explain the meaning of *strong* in terms of ionisation.

\_\_\_\_\_

(1)

- (ii) A student made some ammonium sulfate in a school laboratory.

The student carried out a titration, using a suitable indicator, to find the volumes of sulfuric acid and ammonia solution that should be reacted together.


Name a suitable indicator for strong acid-weak alkali titrations.

\_\_\_\_\_

(1)

- (iii) The student found that 25.0 cm<sup>3</sup> of ammonia solution reacted completely with 32.0 cm<sup>3</sup> of sulfuric acid of concentration 0.050 moles per cubic decimetre.

The equation that represents this reaction is:



Calculate the concentration of this ammonia solution in moles per cubic decimetre.

---

---

---

---

---

Concentration = \_\_\_\_\_ moles per cubic decimetre

(3)

- (iv) Use your answer to (b)(iii) to calculate the concentration of ammonia in grams per cubic decimetre.

(If you did not answer part (b)(iii), assume that the concentration of the ammonia solution is 0.15 moles per cubic decimetre. This is **not** the correct answer to part (b)(iii).)

Relative formula mass of ammonia (NH<sub>3</sub>) = 17.

---

---

---


Concentration = \_\_\_\_\_ grams per cubic decimetre

(2)

(Total 11 marks)

### Q53.

Firework rockets contain fuel and potassium nitrate.





The potassium nitrate provides oxygen for the fuel to react.

- (a) The table shows how a student worked out the relative formula mass ( $M_r$ ) of potassium nitrate.

Some of the numbers are missing.

Relative atomic masses ( $A_r$ ): N = 14; O = 16; K = 39.

Name of atom (symbol)	Number of atoms	$A_r$	Mass
potassium (K)	1	39	39
nitrogen (N)	1	14	14
oxygen (O)		16	
The $M_r$ of potassium nitrate =			101

- (i) The mass of oxygen is not shown in the table.

Draw a ring around the correct mass of oxygen.

**16**                      **32**                      **48**

(1)

- (ii) Draw a ring around the number of oxygen atoms in the formula of potassium nitrate.

**1**                      **2**                      **3**

(1)

- (b) When the fuel reacts with the oxygen an *exothermic* reaction takes place.

What does *exothermic* mean?

---

---

---

---

(2)

- (c) The fuel contains carbon. Carbon reacts with oxygen to make carbon dioxide.

Which **two** statements in the table explain why carbon dioxide is a gas at room temperature?

Tick (✓) the **two** statements.


Statement	Tick (✓)
It has a giant structure	
It has a low boiling point.	
It is made of small molecules.	
It is made of ions.	

(2)  
(Total 6 marks)

**Q54.**

- (a) The table gives information about two isotopes of hydrogen, hydrogen-1 and hydrogen-2.

	Hydrogen-1	Hydrogen-2
Atomic number	1	1
Mass number	1	2



An atom of hydrogen-1 is represented as:

Show how an atom of hydrogen-2 is represented.

(1)

- (b) (i) Calculate the relative formula mass ( $M_r$ ) of water,  $\text{H}_2\text{O}$

Relative atomic masses: H = 1; O = 16.

---



---

Relative formula mass ( $M_r$ ) = \_\_\_\_\_

(1)

- (ii) Simple molecules like water have low boiling points.

Explain why, in terms of molecules.

---



---

(2)

- (c) Molecules of heavy water contain two atoms of hydrogen-2 instead of two atoms of hydrogen-1.

Explain why a molecule of heavy water has more mass than a normal water molecule.

You should refer to the particles in the nucleus of the two different hydrogen atoms in your answer.

---

---

---

---


---

(2)

(Total 6 marks)

### Q55.

Air bags are used to protect the passengers in a car during an accident. When the crash sensor detects an impact it causes a mixture of chemicals to be heated to a high temperature. Reactions take place which produce nitrogen gas. The nitrogen fills the air bag.



- (a) The mixture of chemicals contains sodium azide ( $\text{NaN}_3$ ) which decomposes on heating to form sodium and nitrogen.



A typical air bag contains 130 g of sodium azide.

- (i) Calculate the mass of nitrogen that would be produced when 130 g of sodium azide decomposes.

Relative atomic masses ( $A_r$ ): N = 14; Na = 23

---

---

---

---

Mass of nitrogen = \_\_\_\_\_ g

(3)

- (ii) 1 g of nitrogen has a volume of 0.86 litres at room temperature and pressure.

What volume of nitrogen would be produced from 130 g of sodium azide?

(If you did not answer part (a)(i), assume that the mass of nitrogen produced from 130 g of sodium azide is 80 g. This is **not** the correct answer to part (a)(i).)


---

Volume = \_\_\_\_\_ litres

(1)

- (b) The sodium produced when the sodium azide decomposes is dangerous. The mixture of chemicals contains potassium nitrate and silicon dioxide which help to make the sodium safe.

- (i) Sodium reacts with potassium nitrate to make sodium oxide, potassium oxide and nitrogen. Complete the balancing of the equation for this reaction.



(1)

- (ii) The silicon dioxide reacts with the sodium oxide and potassium oxide to form silicates.

Suggest why sodium oxide and potassium oxide are dangerous in contact with the skin.

---

---

(1)

(Total 6 marks)

