



Knowledge organiser




Science: Foundation Tier

RAYNES
PARK HIGH SCHOOL



cytoplasm	<i>site of chemical reactions in the cell</i>	gel like substance containing enzymes to catalyse the reactions
nucleus	<i>contains genetic material</i>	controls the activities of the cell and codes for proteins
cell membrane	<i>semi permeable</i>	controls the movement of substances in and out of the cell
ribosome	<i>site of protein synthesis</i>	mRNA is translated to an amino acid chain
mitochondrion	<i>site of respiration</i>	where energy is released for the cell to function


animal cell



cell membrane	<i>site of chemical reactions in the cell</i>	gel like substance containing enzymes to catalyse the reactions
bacterial DNA	<i>not in nucleus floats in the cytoplasm</i>	controls the function of the cell
cell wall	NOT made of cellulose	supports and strengthens the cell
plasmid	<i>small rings of DNA</i>	contain additional genes
cytoplasm	<i>semi permeable</i>	controls the movement of substances in and out of the cell

Bacterial cells are much smaller than plant and animal cells


plant cell



contains all the parts of animal cells plus extras

permanent vacuole	<i>contains cell sap</i>	keeps cell turgid, contains sugars and salts in solution
cell wall	<i>made of cellulose</i>	supports and strengthens the cell
chloroplast	<i>site of photosynthesis</i>	contains chlorophyll, absorbs light energy

how a cell changes and becomes specialised
Undifferentiated cells are called **STEM** cells



AQA Cell Structure


Prokaryotes simpler organisms

Specialised cells

specialised animal cells	nerve		<i>carry electrical signals</i>	long branched connections and insulating sheath
	sperm		<i>fertilise an egg</i>	streamlined with a long tail acrosome containing enzymes large number of mitochondria
	muscle		<i>contract to allow movement</i>	contains a large number of mitochondria long
specialised plant cells	root hair		<i>absorb water and minerals from soil</i>	hair like projections to increase the surface area
	xylem		<i>carry water and minerals</i>	TRANSPIRATION - dead cells cell walls toughened by lignin flows in one direction
	phloem		<i>carry glucose</i>	TRANSLOCATION - living cells cells have end plates with holes flows in both directions

Microscopy


$$\text{magnification} = \frac{\text{size of image}}{\text{real size of the object}}$$



Feature	Light (optical) microscope	Electron microscope
Radiation used	Light rays	Electron beams
Max magnification	~ 1500 times	~ 2 000 000 times
Resolution	200nm	0.2nm
Size of microscope	Small and portable	Very large and not portable
Cost	~£100 for a school one	Several £100,000 to £1 million plus

PREFIXES		
Prefix	Multiple	Standard form
centi (cm)	1 cm = 0.01 m	$\times 10^{-2}$
milli (mm)	1 mm = 0.001 m	$\times 10^{-3}$
micro (µm)	1 µm = 0.000 001 m	$\times 10^{-6}$
nano (nm)	1nm = 0.000 000 001 m	$\times 10^{-9}$

largest
↑
smallest




Small intestines	<i>Villi – increase surface area, Good blood supply – to maintain concentration gradient, Thin membranes – short diffusion distance.</i>
Lungs	<i>Alveoli– increase surface area, Good blood supply – to maintain concentration gradient, Thin membranes – short diffusion distance.</i>
Gills in fish	<i>Gill filaments and lamella – increase surface area, Good blood supply – to maintain concentration gradient, Thin membranes – short diffusion distance.</i>
Roots	<i>Root hair cells - increase surface area.</i>
Leaves	<i>Large surface area, thin leaves for short diffusion path, stomata on the lower surface to let O₂ and CO₂ in and out.</i>

ADAPTATIONS FOR DIFFUSION – The greater the difference in concentrations the faster the rate of diffusion.

Cells divide in a series of stages. The genetic material is doubled and then divided into two identical cells.

MITOSIS AND THE CELL CYCLE

Stage 1	Growth	Increase the number of sub-cellular structures e.g. ribosomes and mitochondria.
Stage 2	DNA Synthesis	DNA replicates to form two copies of each chromosome.
Stage 3	Mitosis	One set of chromosomes is pulled to each end of the cell and the nucleus divides. Then the cytoplasm and cell membranes divide to form two cells that are identical to the parent cell.



Mitosis occurs during growth, repair, replacement of cells. Asexual reproduction occurs by mitosis in both plants & simple animals.

AQA Cell Biology 2

Cell division

STEM CELLS

Undifferentiated cell of an organism

Divides to form more cells of the same type, and can differentiate to form many other cell types.

Transport in cells

Diffusion <i>No</i> energy required	<i>Movement of particles in a solution or gas from a higher to a lower concentration</i>	E.g. O ₂ and CO ₂ in gas exchange, urea in kidneys. Factors that affect the rate are concentration, temperature and surface area.
Osmosis <i>No</i> energy required	<i>Movement of water from a dilute solution to a more concentrated solution</i>	E.g. Plants absorb water from the soil by osmosis through their root hair cells. Plants use water for several vital processes including photosynthesis and transporting minerals.
Active transport ENERGY required	<i>Movement of particles from a dilute solution to a more concentrated solution</i>	E.g. movement of mineral ions into roots of plants and the movement of glucose into the small intestines.


Human Embryonic stem cells	<i>Can be cloned and made to differentiate into most cell types</i>	Therapeutic cloning uses same genes so the body does not reject the tissue. Can be a risk of infection
Adult bone marrow stem cells	<i>Can form many types of human cells e.g. blood cells</i>	Tissue is matched to avoid rejection, risk of infection. Only a few types of cells can be formed.
Meristems (plants)	<i>Can differentiate into any plant cell type throughout the life of the plant.</i>	Used to produce clones quickly and economically, e.g. rare species, crop plants with pest /disease resistance

Treatment with stem cells may be able to help conditions such as diabetes and paralysis. Some people object to the use of stem cells on ethical or religious grounds

Enzymes catalyse (increase the rate of) specific reactions in living organisms

An organ system in which organs work together to digest and absorb food.


The 'lock and key theory' is a simplified model to explain enzyme action



Enzymes catalyse specific reactions in living organisms due to the shape of their active site

Digestive enzymes speed up the conversion of large insoluble molecules (food) into small soluble molecules that can be absorbed into the bloodstream

The activity of enzymes is affected by changes in temperature and pH



Enzymes in digestion

The human digestive system

AQA GCSE ORGANISATION Part 1

Principles of organisation

Large changes in temperature or pH can stop the enzyme from working (denature)


<i>Temperature too high</i>	<i>pH too high or too low</i>
Enzyme changes shape (denatures) the substrate no longer fits the active site.	

Non-communicable diseases

More energy consumed in food and drink than used
obesity
Linked to increased rates of cardiovascular disease and development of diabetes type 2.

Food tests

Sugars (glucose)	<i>Benedict's test</i>	Orange to brick red precipitate.
Starch	<i>Iodine test</i>	Turns black.
Biuret	<i>Biuret reagent</i>	Mauve or purple solution.




Carbohydrases (e.g. amylase)		<i>Made in salivary glands, pancreas, small intestine</i>	Break down carbohydrates to simple sugar (e.g. amylase breaks down starch to glucose).
Proteases		<i>Made in stomach, pancreas</i>	Break down protein to amino acids.
Lipases		<i>Made in pancreas (works in small intestine)</i>	Break down lipids (fats) to glycerol and fatty acids.
Bile (not an enzyme)		<i>Made in liver, stored in gall bladder.</i>	Emulsifies lipids to increase surface area to increase the rate of lipid break down by lipase. Changes pH to neutral for lipase to work

The products of digestion are used to build new carbohydrates, lipids and proteins. Some glucose is used for respiration.


Cells, tissues, organs and systems

Cells		<i>e.g. muscle cells</i>	The basic building blocks of all living organisms.
Tissues		<i>e.g. muscle tissue</i>	A group of cells with a similar structure and function.
Organs		<i>e.g. the heart</i>	Aggregations (working together) of tissues performing a specific function.
Organ systems		<i>e.g. the circulatory system</i>	Organs working together to form organ systems, which work together to form an organism.



Blood vessels

The heart is an organ that pumps blood around the body in a double circulatory system



Different structure in the heart have different functions	<i>Right ventricle</i>	Pumps blood to the lungs where gas exchange takes place.
	<i>Left ventricle</i>	Pumps blood around the rest of the body.
	<i>Pacemaker (in the right atrium)</i>	Controls the natural resting heart rate. Artificial electrical pacemakers can be fitted to correct irregularities.
	<i>Coronary arteries</i>	Carry oxygenated blood to the cardiac muscle.
	<i>Heart valves</i>	Prevent blood in the heart from flowing in the wrong direction.

Artery	Vein	Capillary
<i>Carry blood away from the heart</i>	<i>Carry blood to the heart</i>	<i>Connects arteries and veins</i>
Thick muscular walls, small lumen, carry blood under high pressure, carry oxygenated blood (except for the pulmonary artery).	Thin walls, large lumen, carry blood under low pressure, have valves to stop flow in the wrong direction, carry deoxygenated blood (except for the pulmonary vein).	One cell thick to allow diffusion, Carry blood under very low pressure.

Heart

Blood

Blood is a tissue consisting of plasma, in which blood cells, white blood cells and platelets are suspended


AQA GCSE ORGANISATION part 2

Lungs and gas exchange

The heart pumps low oxygen/high carbon dioxide blood to the lungs

Plasma (55%)	<i>Pale yellow fluid</i>	Transports CO ₂ , hormones and waste.
Red blood cells (45%)	<i>Carries oxygen</i>	Large surface area, no nucleus, full of haemoglobin.
White blood cells (<1%)	<i>Part of the immune system</i>	Some produce antibodies, others surround and engulf pathogens.
Platelets (<1%)	<i>Fragments of cells</i>	Clump together to form blood clots.




Trachea	<i>Carries air to/from the lungs</i>	Rings of cartilage protect the airway.
Bronchioles	<i>Carries air to/from the air sacs (alveoli)</i>	Splits into multiple pathways to reach all the air sacs.
Alveoli	<i>Site of gas exchange in the lungs</i>	Maximises surface area for efficient gas exchange.
Capillaries	<i>Allows gas exchange between into/out of blood</i>	Oxygen diffuses into the blood and carbon dioxide diffuses out.



Disease	Cause	Effect	Treatment
Coronary heart disease (CHD)	<i>A build up for fatty substances in the coronary arteries (atherosclerosis)</i>	Oxygen-ated blood cannot get to the cardiac muscle.	Stents: inserted into the blocked artery to open it up. Statins: lower harmful cholesterol.
Faulty heart valves	<i>Valves don't open or close properly</i>	Blood can leak or flow in the wrong direction	Biological valve transplant or a mechanical valve can be inserted

Plant organ systems

The roots, stem and leaves form a plant organ system for transport of substances around the plant



Epidermal tissues	<i>Waxy cuticle (top layer of the leaf)</i>	Reduces water loss from the leaf
	<i>Guard cells and stomata</i>	Guard cells open and close the stomata to control water loss and allow for gas exchange (oxygen and carbon dioxide).
Palisade mesophyll	<i>Palisade cells</i>	Cells near the top surface of the leaf that are packed with chloroplasts that contain chlorophyll. Both adaptations maximize photosynthesis.
Spongy mesophyll	<i>Air spaces in the leaf between cells</i>	Increased surface area for gas exchange so that carbon dioxide can diffuse into photosynthesising cells.
xylem	<i>Hollow tubes strengthened by lignin adapted for the transportation of water in the transpiration stream</i>	Allows transport of water and mineral ions from the roots to the stem and the leaves.
phloem	<i>Cell sap moves from one phloem cell to the next through pores in the end walls</i>	Transports dissolved sugars from the leaves to the rest of the plant for immediate use or storage (translocation).
Meristem tissue	<i>New cells (roots and shoot tips) are made here including root hair cells</i>	Root hair cells have an increased surface area for the uptake of water by osmosis, and mineral ions by active transport.



Cancer
Non-communicable diseases
The result of changes in DNA that lead to uncontrolled growth and division

Benign tumour	Contained in one area of the body (usually by a membrane) – not cancer.
Malignant tumour	Invasively spread to different parts of the body to form secondary tumours.

Some cancers have genetic risk factors. Carcinogens and ionising radiation increase the risk of cancer by changing/ damaging DNA

Risk factors for heart/lung disease and certain types of cancer include drinking alcohol, diet, obesity and smoking





These risks factors can also affect the brain, liver and the health of unborn babies




A potometer is used to measure the amount of water lost over time (rate of transpiration)

Transpiration


Transpiration
The rate at which water is lost from the leaves of a plant. The transpiration stream is the column of water moving through the roots, stem and leaves
Temperature, humidity, air movement and light intensity affect the rate of transpiration.



The shape of the graph for light intensity is the same for temperature (energy)



Phagocytes	Phagocytosis	Phagocytes engulf the pathogens and digest them.
Lymphocytes	Antibody production	Specific antibodies destroy the pathogen. This takes time so an infection can occur. If a person is infected again by the same pathogen, the lymphocytes make antibodies much faster.
	Antitoxin production	Antitoxin is a type of antibody produced to counteract the toxins produced by bacteria.



Pathogens are identified by white blood cells by the different proteins on their surfaces **ANTIGENS**.

White blood cells are part of the immune system

Immune system

Non-specific defence systems

The human body has several non specific ways of defending itself from pathogens getting in

	Nose	Nasal hairs, sticky mucus and cilia prevent pathogens entering through the nostrils.
	Trachea and bronchus (respiratory system)	Lined with mucus to trap dust and pathogens. Cilia move the mucus upwards to be swallowed.
	Stomach acid	Stomach acid (pH1) kills most ingested pathogens.
	Skin	Hard to penetrate waterproof barrier. Glands secrete oil which kill microbes

Detection and identification of plant diseases (bio only)

Detection	Identification
<i>Stunted growth</i>	Reference using gardening manual or website, laboratory test for pathogens, testing kit using monoclonal antibodies.
<i>Spots on leaves</i>	
<i>Area of decay</i>	
<i>growths</i>	
<i>Malformed stem/leaves</i>	
<i>Discolouration</i>	
<i>Presence of pests</i>	

AQA GCSE INFECTION AND RESPONSE part 1

Plants have several ways of defending themselves from pathogens and animals

Physical	Mechanical
Thick waxy layers, cell walls stop pathogen entry	Thorns, curling up leaves to prevent being eaten
Chemical	
Antibacterial and toxins made by plant	

Human defence systems

Pathogens may infect plants or animals and can be spread by direct contact, water or air

Nitrate ions needed for protein synthesis – lack of nitrate = stunted growth.

Magnesium ions needed to make chlorophyll – not enough leads to chlorosis – leaves turn yellow.

Bacteria may produce toxins that damage tissues and make us feel ill

Viruses	Bacteria (prokaryotes)	Protists (eukaryotes)	Fungi (eukaryotes)
<i>e.g. cold, influenza, measles, HIV, tobacco mosaic virus</i>	<i>e.g. tuberculosis (TB), Salmonella, Gonorrhoea</i>	<i>e.g. dysentery, sleeping sickness, malaria</i>	<i>e.g. athlete's foot, thrush, rose black spot</i>
DNA or RNA surrounded by a protein coat	No membrane bound organelles (no chloroplasts, mitochondria or nucleus). Cell wall. Single celled organisms	Membrane bound organelles. Usually single celled.	Membrane bound organelles, cell wall made of chitin. Single celled or multi-cellular

Pathogens are microorganisms that cause infectious disease




Pathogens

Communicable diseases

Viruses live and reproduce inside cells causing damage

Pathogen	Disease	Symptoms	Method of transmission	Control of spread
Virus	Measles	Fever, red skin rash.	Droplet infection from sneezes and coughs.	Vaccination as a child.
Virus	HIV	Initially flu like systems, serious damage to immune system.	Sexual contact and exchange of body fluids.	Anti-retroviral drugs and use of condoms.
Virus	Tobacco mosaic virus	Mosaic pattern on leaves.	Enters via wounds in epidermis caused by pests.	Remove infected leaves and control pests that damage the leaves.
Bacteria	Salmonella	Fever, cramp, vomiting, diarrhoea.	Food prepared in unhygienic conditions or not cooked properly.	Improve food hygiene, wash hands, vaccinate poultry, cook food thoroughly.
Bacteria	Gonorrhoea	Green discharge from penis or vagina.	Direct sexual contact or exchange of body fluids.	Use condoms. Treatment using antibiotics.
Protists	Malaria	Recurrent fever.	By an animal vector (mosquitoes).	Prevent breeding of mosquitoes. Use of nets to prevent bites.
Fungus	Rose black spot	Purple black spots on leaves.	Spores carried via wind or water.	Remove infected leaves. Spray with fungicide.

Most new drugs are synthesised by chemists in the pharmaceutical industry.

Traditionally drugs were extracted from plants and microorganisms		
<i>Digitalis</i>	<i>Aspirin</i>	<i>Penicillin</i>
Extracted from foxglove plants and used as a heart drug	A painkiller and anti-inflammatory that was first found in willow bark	Discovered by Alexander Fleming from the <i>Penicillium</i> mould and used as an antibiotic
		

Drugs have to be tested and trialled before to check they are safe and effective

Antibiotics and painkillers

Bacteria can mutate

Sometimes this makes them resistant to antibiotic drugs.

Antibiotics have greatly reduced deaths from infectious bacterial disease

antibiotics	<i>e.g. penicillin</i>	Kill infective bacteria inside the body. Specific bacterial infections require specific antibiotics.
Painkillers and other medicines	<i>e.g. aspirin, paracetamol, ibuprofen</i>	Drugs that are used to treat the symptoms of a disease. They do not kill pathogens

Antibiotics cannot be used to treat viral pathogens

Discovery and drug development

AQA INFECTION AND RESPONSE

Vaccination

It is difficult to develop drugs to kill viruses without harming body tissues because viruses live and reproduce inside cells

New drugs are extensively tested for:	Efficacy	Make sure the drug works
	Toxicity	Check that the drug is not poisonous
	Dose	The most suitable amount to take



Used to immunise a large proportion of the population to prevent the spread of a pathogen

Preclinical trials - using cells, tissues and live animals - must be carried out before the drug can be tested on humans.

Clinical trials use healthy volunteers and patients



Double blind trial: patients and scientists do not know who receives the new drug or placebo until the end of the trial. This avoids bias.

Vaccination	<i>Small amount of dead or inactive form of the pathogen</i>	<i>1st infection by pathogen</i>	White blood cells detect pathogens in the vaccine. Antibodies are released into the blood.
		<i>Re-infection by the same pathogen</i>	White blood cells detect pathogens. Antibodies are made much faster and in larger amounts.

<i>Stage 1</i>	<i>Stage 2</i>	<i>Stage 3</i>	<i>Stage 4</i>
Healthy volunteers try small dose of the drug to check it is safe record any side effects	A small number of patients try the drug at a low dose to see if it works	A larger number of patients; different doses are trialled to find the optimum dose	A double blind trial will occur. The patients are divided into groups. Some will be given the drug and some a placebo.

A placebo can look identical to the new drug but contain no active ingredients

A person is unlikely to suffer the symptoms of the harmful disease and it's spread in a population is prevented




Respiration, stored as insoluble starch, fats or oils for storage, cellulose for cell walls, combine with nitrates from the soil to form amino acids for protein synthesis

Plants use the glucose produced in photosynthesis in a variety of ways

Photosynthetic reaction

The plant manufactures glucose from carbon dioxide and water using energy transferred from the environment to the chloroplasts by light




The rate of photosynthesis is affected by temperature, light intensity, carbon dioxide concentration, and the amount of chlorophyll

Factor	How the rate is affected	Limiting factors (why the rate stops going up)
Temperature	<i>As the temperature of the environment the plant is in increases rate of photosynthesis increases (up to a point) as there is more energy for the chemical reaction.</i>	Photosynthesis is an enzyme controlled reaction. If the temperature increases too much, then the enzymes become denatured and the rate of reaction will decrease and stop
Light intensity	<i>Light intensity increases as the distance between the plant and the light sources increases. As light intensity increases so does the rate of photosynthesis (up to a point) as more energy is available for the chemical reaction.</i>	At point X another factor is limiting the rate of photosynthesis. This could be carbon dioxide concentration, temperature or the amount of chlorophyll
Carbon dioxide concentration	<i>Carbon dioxide is needed for plants to make glucose. The rate of photosynthesis will increase when a plant is given higher concentrations of carbon dioxide (up to a point).</i>	At point X another factor is limiting the rate of photosynthesis. This could be light intensity, temperature or the amount of chlorophyll
Amount of chlorophyll	<i>Chlorophyll is a photosynthetic pigment that absorbs light and allows the reaction between water and carbon dioxide to occur (photosynthesis)</i>	Another factor could limit the rate of photosynthesis. This could be light intensity, temperature or the carbon dioxide concentration

AQA GCSE BIOENERGETICS part 1


Rate of photosynthesis

Rate of photosynthesis




Graph lines C and D: If temperature is increased by 10°C then a slight increase in rate of photosynthesis occurs.

Explain graphs of two or three factors and decide which is the limiting factor




Graph lines A and D: If carbon dioxide concentration and temperature are increased the rate of photosynthesis increases significantly up to a point.




Graph Lines A and B: If carbon dioxide concentration is increased from 0.01% to 0.1% then a large increase in rate occurs up to a point.




Graph line A: Rate could be limited by temperature and/or amount of chlorophyll. Plant tissue can be damaged when carbon dioxide concentrations exceed 0.1%



During long periods of vigorous activity muscles become fatigued and stop contracting efficiently

An organism will receive all the energy it needs for living processes as a result of the energy transferred from respiration	<i>For movement</i>	 Smooth muscle cells	To enable muscles to contract in animals.
	<i>For keeping warm</i>		To keep a steady body temperature in a cold environment.
	<i>For chemical reactions</i>		To build larger molecules from smaller one.



Electron micrograph of a mitochondrion

Response to exercise

During exercise the human body reacts to increased demand for energy	<i>Heart rate increases</i>	Top pump oxygenated blood faster to the muscle tissues and cells.
	<i>Breathing rate and breath volume increase</i>	This increases the amount of oxygen entering the blood stream.

Respiration
AQA GCSE BIOENERGETICS part 2



Cellular respiration is an exothermic reaction which is continuously occurring in all living cells

Metabolism is the sum of all the reactions in a cell or the body

Metabolism	<i>The energy transferred by respiration in cells is used by the organism for the continual enzyme controlled processes of metabolism.</i>	Conversion of glucose to starch, glycogen and cellulose.
		The formation of lipid molecules from a molecule of glycerol and three molecules of fatty acid.
		The use of glucose and nitrate ions to form amino acids which in turn are used to synthesise proteins.
		Respiration
		Breakdown of excess proteins to form urea for excretion.

Anaerobic respiration in plant and yeast cells
The end products are ethanol and carbon dioxide. Anaerobic respiration in yeast cells is called fermentation
glucose → ethanol + carbon dioxide

This process is economically important in the manufacture of alcoholic drinks and bread.



Anaerobic respiration
Respiration when oxygen is in short supply. Occurs during intensive exercise
During hard exercise, muscle cells are respiring so fast that blood cannot transport enough oxygen to meet their needs.
Glucose is partially oxidised to produce lactic acid which builds up in muscle tissue causing them to become painful and fatigued.
glucose → lactic acid

Anaerobic respiration releases a much smaller amount of energy than aerobic respiration.

The incomplete oxidation of glucose causes a build up of lactic acid and creates an oxygen debt

Aerobic respiration
Respiration with oxygen. Occurs inside the mitochondria continuously
Glucose is oxidised by oxygen to transfer the energy the organism needs to perform its functions.
$$C_6H_{12}O_6 + O_2 \rightarrow CO_2 + H_2O$$

glucose + oxygen → carbon dioxide + water


Aerobic respiration releases a large amount of energy from each glucose molecule

Human control systems include	Cells called receptors	Detect stimuli (changes in environment).
	Coordination centres	e.g. brain, spinal cord and pancreas that receive information from receptors.
	Effectors	Muscles or glands, which bring about responses to restore optimum levels.

Enables humans to react to their surroundings and to co-ordinate their behaviour

AQA GCSE HOMEOSTASIS AND RESPONSE part 1


The human nervous system



Information from receptors passes along cells (neurones) as electrical impulses to the central nervous system (CNS)



The CNS is the brain and the spinal cord.

Coordinates the response of effectors; muscles contracting or glands secreting hormones




Typical motor neurone

Synapse (gap where two neurones meet).



Reflex arc	Receptor	Detect stimuli.
	Sensory neurone	Long axon carries impulse from receptor to spinal cord.
	Synapse	Gap where neurones meet. Chemical message using neurotransmitter.
	Relay neurone	Allows impulses to travel between sensory and motor neurones in the spinal cord.
	Motor neurone	Long axon carries impulse from receptor to effector.
	Effector	Muscle or gland that carries out response.



Reflex actions are automatic and rapid; they do not involve the conscious part of the brain and can protect humans from harm.

Response to internal and external change

Controls in the human body	<i>Blood glucose concentration</i>	These automatic control systems may involve nervous responses or chemical responses.
	<i>Body temperature</i>	
	<i>Water levels</i>	


The regulation of internal conditions of a cell or organism to maintain optimum conditions for function.

Homeostasis maintains optimal conditions for enzyme action and all cell functions.

Homeostasis

AQA GCSE HOMEOSTASIS AND RESPONSE PART 2

Human endocrine system



Endocrine system	<i>Composed of glands which secrete chemicals called hormones directly into the bloodstream.</i>	The blood carries the hormone to a target organ where it produces an effect. Compared to the nervous system effects are slower but act for longer.
-------------------------	--	--

Pituitary gland	<i>'Master gland'; secretes several hormones into the blood</i>	Stimulates other glands to produce hormones to bring about effects.
------------------------	---	---

Control of blood glucose concentration

Blood glucose concentration	
<i>Monitored and controlled by the pancreas</i>	
Too high	(HT only) Too low
Pancreas produces the hormone insulin, glucose moves from the blood into the cells. In liver and muscle cells excess glucose is converted to glycogen for storage.	Pancreas produces the hormone glucagon that causes glycogen to be converted into glucose and released into the blood.

Negative feedback (HT only)	<i>Adrenaline</i>	Produced in adrenal glands, increases breathing/heart rate, blood flow to muscles, conversion glycogen to glucose. Prepares body for 'fight or flight'.
	<i>Thyroxine</i>	Produced in the thyroid gland, stimulates the basal metabolic rate. Important in growth and development.

Increasing thyroxine levels prevent the release of thyroid stimulating hormone which stops the release of thyroxine.

Diabetes	
<i>Type 1</i>	<i>Type 2</i>
Pancreas fails to produce sufficient insulin leading to uncontrolled blood glucose levels. Normally treated by insulin injection.	Obesity is a risk factor. Body cells no longer respond to insulin. Common treatments include changing by diet and increasing exercise.


(HT) Rising glucose levels inhibit the release of glucagon in a negative feedback system. Insulin is released to reduce glucose levels and which cause the pancreas to release glucagon

**AQA GCSE
HOMEOSTASIS AND
RESPONSE PART 3**

**Hormones in human
reproduction**


Contraception

During puberty reproductive hormones cause secondary sexual characteristics to develop	
<i>Oestrogen (main female reproductive hormone)</i>	<i>Testosterone (main male reproductive hormone)</i>
Produced in the ovaries. At puberty eggs begin to mature releasing one every 28 days – ovulation.	Produced in the testes stimulating sperm production.




Fertility can be controlled by hormonal and non hormonal methods	<i>Oral contraceptives</i>	Contain hormones to inhibit FSH production so that no eggs mature.
	<i>Injection, implant, skin patch</i>	For slow release of progesterone to inhibit the maturation and release of eggs for months or years.
	<i>Barrier methods</i>	Condoms or diaphragms which prevent sperm reaching the egg.
	<i>Intrauterine devices</i>	Prevent implantation of an embryo or release a hormone.
	<i>Spermicidal agents</i>	Kill or disable sperm.
	<i>Abstaining</i>	Avoiding intercourse when an egg may be in the oviduct.
	<i>Surgery</i>	Male or female sterilisation.

Menstrual cycle	<i>Follicle stimulating hormone (FSH)</i>	Causes maturation of an egg in the ovary.	(HT) FSH stimulates ovaries to produce oestrogen.
	<i>Luteinising hormone (LH)</i>	Stimulates release of an egg.	(HT) Oestrogen stops FSH production and stimulates LH production in pituitary gland.
	<i>Oestrogen and progesterone</i>	Maintain uterus lining.	



Meiosis halves the number of chromosomes

<p>Gametes are made in reproductive organs (in animals ovaries and testes)</p>	<p><i>Cells divide by meiosis to form gametes</i></p>	<p>Copies of the genetic information are made.</p>
		<p>The cell divides twice to form four gametes each with single set of chromosomes.</p>
		<p>All gametes are genetically different from each other.</p>



<p>Sexual reproduction involves the fusion of male and female gametes.</p>	<p><i>Sperm and egg in animals.</i></p> <p><i>Pollen and egg cells in flowering plants.</i></p>	<p>Produced by meiosis. There is mixing of genetic information which leads to a variety in the offspring.</p>
<p>Asexual reproduction involves only one parent and no fusion of gametes.</p>	<p><i>e.g. cloning of females only in an aphid population.</i></p>	<p>Only mitosis is involved. There is no mixing of genetic information. This leads to genetically identical clones.</p>

Gametes join at fertilisation to restore the number of chromosomes

The new cell divides by mitosis. The number of cells increase. As the embryo develops cells differentiate.

Meiosis

Meiosis leads to non-identical cells being formed while mitosis leads to identical cells being formed

DNA and the genome


Sexual and asexual reproduction

Genetic material in the nucleus is composed of a chemical called DNA.



AQA GCSE INHERITANCE, VARIATION AND EVOLUTION Part 1

<p>DNA structure</p>
<p><i>Polymer made up of two strands forming a double helix.</i></p>
<p>Contained in structures called chromosomes. A gene is a small section of DNA on a chromosome. Each gene codes for a sequence of amino acids to make a specific protein.</p>



The genome is the entire genetic material of an organism.

<p>The whole human genome has now been studied.</p>	<p><i>It is of great importance for future medical developments</i></p>	<p>Searching for genes linked to different types of disease.</p>
		<p>Understanding and treatment of inherited disorders.</p>
		<p>Tracing migration patterns from the past.</p>

Embryo screening: small piece of developing placenta removed to check for presence of faulty genes

Gene therapy: replacing the faulty allele in somatic cells with a normal allele

Very rarely a mutation will lead to a new phenotype which if is suited to environmental change can lead to rapid change in the species.

Embryo screening /gene therapy issues	Economic	Costly and not 100% reliable.
	Social	Not available to everyone (due to cost).
	Ethical	Should only 'healthy' embryos be implanted following screening.

Embryo screening and gene therapy may alleviate suffering

Some disorders are inherited. They are caused by the inheritance of certain alleles

Ordinary human body cells contain 23 pairs of chromosomes

One pair of chromosomes carry the genes that determine sex


	Female	Male
	XX	XY
Gametes	X	Y
X	XX	XY
X	XX	XY

The probability of a male of female child is 50%. The ratio is 1:1

Variation: difference in the characteristics of individuals in a population may be due to

Genetic causes (inheritance)	There is usually extensive genetic variation within the population of a species e.g. hair colour, skin colour, height that can also be affected by environment e.g. nutrition, sunlight.
Environmental causes (condition they have developed in)	
A combination of genes and environment	

Using a family tree: if the father was homozygous dominant then all of the offspring would have the disorder. He must be heterozygous




Inherited disorders

Polydactyly	Cystic fibrosis
Caused by inheriting a dominant allele.	Caused by inheriting a recessive allele (both parents have to at least carry it).
Causes a person/animal to have extra toes or fingers.	A disorder of the cell membrane. Patients cannot control the viscosity of their mucus.

Sex determination

Using a punnet square (using mouse fur colour as an example)

Parent phenotype	Black fur	White fur
Parent genotype	BB	bb
What gametes are present	In each egg B	In each sperm b



The probability of black fur offspring phenotype is 100%. All offspring genotypes are heterozygous (Bb).

Crossing two heterozygous mice (Bb)

Gametes	B	b
B	BB	Bb
b	Bb	bb

The probability of black fur is 75% and white fur 25%. The ratio of black to white mice is 3:1

All genetic variation arises in mutation, most have no effect on phenotype, some influence but very few determine phenotype.

Variation

The genome and its interaction with the environment influence the development of phenotypes

AQA GCSE INHERITANCE, VARIATION AND EVOLUTION PART 2

Define terms linked to genetics

Gamete	Sex cells produced in meiosis.
Chromosome	A long chain of DNA found in the nucleus.
Gene	Small section of DNA that codes for a particular protein.
Allele	Alternate forms of the same gene.
Dominant	A type of allele – always expressed if only one copy present and when paired with a recessive allele.
Recessive	A type of allele – only expressed when paired with another recessive allele.
Homozygous	Pair of the same alleles, dominant or recessive.
Heterozygous	Two different alleles are present 1 dominant and 1 recessive.
Genotype	Alleles that are present for a particular feature e.g. Bb or bb
Phenotype	Physical expression of an allele combination e.g. black fur, blonde hair, blue eyes.

Some characteristics are controlled by a single gene e.g. fur colour, colour blindness.

The alleles present, or genotype operate at a molecular level to develop characteristics that can be expressed as a phenotype.

Most characteristics are as a result of multiple genes interacting.

Genetic inheritance

The concept of probability in predicting results of a single gene cross.

Dominant and recessive allele combinations

Dominant	Recessive
Represented by a capital letter e.g. B.	Represented by a lower case letter e.g. b.

3 possible combinations:
Homozygous dominant BB
Heterozygous dominant Bb
Homozygous recessive bb

Over time this results in the formation of new species.

The theory of evolution by natural selection.

Species of all living things have evolved from simple life forms that first developed 3 billion years ago.

Through natural selection of variants (genotypes) that give rise to phenotypes best suited to their environment or environmental change e.g. stronger, faster. This allows for variants to pass on their genotype to the next generation.




If two populations of one species become so different in phenotype that they can no longer interbreed to produce fertile offspring they have formed two new species.



Classification of living organisms

Evolutionary trees are a method used by scientists to show how organisms are related

Use current classification data for living organisms and fossil data for extinct organisms



Choosing characteristics

Desired characteristics are chosen for usefulness or appearance

Disease resistance in food crops.



Animals which produce more meat or milk.



Domestic dogs with a gentle nature.



Large or unusual flowers.



Selective breeding can lead to 'inbreeding' where some breeds are particularly prone to disease or inherited defects e.g. British Bulldogs have breathing difficulties.

Humans have been doing this for thousands of years since they first bred food from crops and domesticated animals.

A change in the inherited characteristics of a population over time through the process of natural selection.

Evolution

AQA GCSE INHERITANCE VARIATION AND EVOLUTION PART 3

The process by which humans breed plants/animals for particular genetic characteristics

Selective breeding

Selective breeding

Choosing parents with the desired characteristics from a mixed population

Chosen parents are bred together.



From the offspring those with desired characteristics are bred together.



Repeat over several generations until all the offspring show the desired characteristics.

AQA GCSE INHERITANCE VARIATION AND EVOLUTION PART 4

Classification of living organisms

The full human classification

Carl Linnaeus classified living things	Kingdom	Animalia
	Phylum	Chordata
	Class	Mammalia
	Order	Primates
	Family	Hominidae
	Genus	<i>Homo</i>
	Species	<i>sapiens</i>

Due to improvements in microscopes, and the understanding of biochemical processes, new models of classification were proposed.

Carl Woese
3 domain based on chemical analysis.
Archaea (primitive bacteria), true bacteria, eukaryota.

Organisms are named by the binomial system of genus and species. Humans are *Homo sapiens*

Evidence for evolution

Fossils and antibiotic resistance in bacteria provide evidence for evolution.

Antibiotic resistant bacteria	Mutations produce antibiotic resistant strains which can spread	Resistant strains are not killed.
		Strain survives and reproduces.
		People have no immunity to strain and treatment is ineffective.

Extinction
When no members of a species survive
Due to extreme geological events, disease, climate change, habitat destruction, hunting by humans.



Fossils tell scientists how much or how little different organisms have changed over time.


Fossils
'remains' of ancient organisms which are found in rocks
Parts of organism that have not decayed as necessary conditions are absent.
Parts of the organism replaced by minerals as they decay.
Preserved traces of organisms such as footprints, burrows and rootlet traces.

Early forms of life were soft bodied and few traces are left behind and have been destroyed by geological activity, cannot be certain about how life began.

Evolution is widely accepted. Evidence is now available as it has been shown that characteristics are passed on to offspring in genes.

Farmers optimise conditions for making compost for use as a natural fertiliser.

Anaerobic decay in biogas generators produces methane gas, used as a fuel.




Factors affecting rate of decay
Temperature, water, oxygen
Increase the rate of decay. In enzyme controlled reactions raising the temperature too high will denature the enzymes.

Breakdown of dead organisms releases mineral ions can into the soil.

Ecosystem	Environment	The conditions surrounding an organism; abiotic and biotic.
	Habitat	Place where organisms live e.g. woodland, lake.
	Population	Individuals of a species living in a habitat.
	Community	Populations of different species living in a habitat.





Organisms require a supply of materials from their surroundings and from the other living organisms.

Bacteria respire when breaking down dead organisms releasing CO₂.



Materials are recycled to provide the building blocks for future organisms

Photosynthetic organisms are the producers of biomass for life on Earth

Food chains			
Feeding relationships in a community			
Producer	Primary consumer	Secondary consumer	Tertiary consumer
			
Grass → Grasshopper → Mouse → Owl			
All food chains begin with a producer e.g. grass that is usually a green plant or photosynthetic algae.		Consumers that kill and eat other animals are predators and those eaten are prey.	

Surviving and reproducing	Competition	Plants in a community or habitat compete with each other for light, space, water and mineral ions. Animals compete with each other for food, mates and territory.
	Interdependence	Species depend on each other for food, shelter, pollination, seed dispersal etc. Removing a species can affect the whole community

EXAMPLE: Introduction of grey squirrels to UK increased competition for food for red squirrels. The greys also carry a pathogen that kills reds.

EXAMPLE: climate change is leading to more dissolved CO₂ in oceans lowering the pH of the water affecting organisms living there.

Decomposition and material cycling

Interdependence and competition

AQA GCSE ECOLOGY PART 1


Levels of organisation

Organisms adaptations enable them to survive in conditions where they normally live.




Abiotic and biotic factors.

Abiotic	Biotic
<i>Non-living factors that affect a community</i>	<i>Living factors that affect a community</i>
Living intensity.	Availability of food.
Temperature.	New predators arriving.
Moisture levels.	
Soil pH, mineral content.	New pathogens.
Wind intensity and direction.	
Carbon dioxide levels for a plant.	One species outcompeting so numbers are no longer sufficient to breed
Oxygen levels for aquatic organisms.	

Adaptations may be structural, behavioural or functional.



In a stable community the numbers of predators and prey rise and fall in cycles.

Adaptations		
Plants	Animals	Extremophiles
Cactus in dry, hot desert	Polar bear in extreme cold artic	Deep sea vent bacteria
		
No leaves to reduce water loss, wide deep roots for absorbing water.	Hollow hairs to trap layer of heat. Thick layer of fat for insulation.	Populations form in thick layers to protect outer layers from extreme heat of vent.


Global warming

Levels of CO₂ and methane in the atmosphere are increasing.

Decreased land availability from sea level rise, temperature rise damages delicate habitats, extreme weather events harm populations of plants and animals.

There is a global consensus about global warming and climate change based on systematic reviews of thousands of peer reviewed publications.

Global warming



AQA GCSE ECOLOGY PART 2

Maintaining biodiversity

Human activity can have a positive impact on biodiversity

Scientists and concerned citizens

Put in place programmes to reduce the negative impacts of humans on ecosystems and biodiversity

Breeding programmes for endangered species.

Protection and regeneration of rare habitats.

Reintroduction of field margins and hedgerows in agricultural areas where farmers grow only one type of crop.

Reduction of deforestation and CO₂ emissions by some governments.

Recycling resources rather than dumping waste in landfill.

Some of the programmes potentially conflict with human needs for land use, food production and high living standards.

Maintain a great biodiversity	Ensures the stability of ecosystems	By reducing the dependence on one species on another for food, shelter, maintenance of the physical environment.
	Future of human species	Many human activities are reduction biodiversity and only recently measures have been taken to stop it.

Human activity can have a negative impact on biodiversity



Pollution kills plants and animals which can reduce biodiversity.

Biodiversity is the variety of all different species of organisms on Earth, or within an ecosystem

Biodiversity

Biodiversity and the effect of human interaction on the ecosystem

Waste management	Rapid growth in human population and higher standard of living	More resources used and more waste produced.
		Pollution in water; sewage, fertiliser or toxic chemicals.
		Pollution in air; smoke or acidic gases.
		Pollution on land; landfill and toxic chemicals.

Experimental methods are used to determine the distribution and abundance of a species.



Sampling techniques	Quadrats	Organisms are counted within a randomly placed square
	Transects	Organisms are counted along a belt (transect) of the ecosystem.



AQA GCSE ECOLOGY PART 3

Waste, land use and deforestation

Land use
Humans reduce the amount of land and habitats available for other plants, animals and microorganisms.
Building and quarrying.
Farming for animals and food crops.
Dumping waste.
Destruction of peat bogs to produce cheap compost for gardeners/farmers to increase food production.

Processing data	
Median	Middle value in a sample.
Mode	Most occurring value in a sample.
Mean	The sum of all the value in a sample divided by the sample number.



Large scale deforestation
In tropical areas (e.g. rain forest) has occurred to:
Provide land for cattle and rice fields, grow crops for biofuels.

The decay or burning of peat release CO₂ into the atmosphere.




This conflicts with conserving peat bogs and peatlands as habitats for biodiversity and reduce CO₂ emissions.

Deforestation reduces biodiversity and removes a sink for increasing the amount CO₂ in the atmosphere.

Atoms, elements and compounds

Atom	<i>The smallest part of an element that can exist</i>	Have a radius of around 0.1 nanometres and have no charge (0).
Element	<i>Contains only one type of atom</i>	Around 100 different elements each one is represented by a symbol e.g. O, Na, Br.
Compound	<i>Two or more elements chemically combined</i>	Compounds can only be separated into elements by chemical reactions.

Pre 1900		<i>Tiny solid spheres that could not be divided</i>	Before the discovery of the electron, John Dalton said the solid sphere made up the different elements.
1897 'plum pudding'		<i>A ball of positive charge with negative electrons embedded in it</i>	JJ Thompson's experiments showed that an atom must contain small negative charges (discovery of electrons).
1909 nuclear model		<i>Positively charge nucleus at the centre surrounded negative electrons</i>	Ernest Rutherford's alpha particle scattering experiment showed that the mass was concentrated at the centre of the atom.
1913 Bohr model		<i>Electrons orbit the nucleus at specific distances</i>	Niels Bohr proposed that electrons orbited in fixed shells; this was supported by experimental observations.



Central nucleus	Contains protons and neutrons
Electron shells	Contains electrons

Electronic shell	Max number of electrons
1	2
2	8
3	8
4	2

Electronic structures

Name of Particle	Relative Charge	Relative Mass
Proton	+1	1
Neutron	0	1
Electron	-1	Very small

Relative electrical charges of subatomic particles

<div style="display: flex; flex-direction: column; align-items: center;"> <div style="margin-bottom: 5px;">7 ←</div> <div style="margin-bottom: 5px;">Li</div> <div>3 ←</div> </div>	Mass number	<i>The sum of the protons and neutrons in the nucleus</i>	
	Atomic number	<i>The number of protons in the atom</i>	Number of electrons = number of protons

AQA GCSE Atomic structure and periodic table part 1

The development of the model of the atom

James Chadwick	<i>Provided the evidence to show the existence of neutrons within the nucleus</i>
-----------------------	---

Rutherford's scattering experiment	<p><i>A beam of alpha particles are directed at a very thin gold foil</i></p>	<p>Most of the alpha particles passed right through. A few (+) alpha particles were deflected by the positive nucleus. A tiny number of particles reflected back from the nucleus.</p>
---	---	--

Mixtures	<i>Two or more elements or compounds not chemically combined together</i>	Can be separated by physical processes.
-----------------	---	---

Chemical equations	<i>Show chemical reactions - need reactant(s) and product(s) energy always involves and energy change</i>	Law of conservation of mass states the total mass of products = the total mass of reactants.
---------------------------	---	--

Word equations	<i>Uses words to show reaction</i> reactants → products magnesium + oxygen → magnesium oxide	Does not show what is happening to the atoms or the number of atoms.
Symbol equations	<i>Uses symbols to show reaction</i> reactants → products $2Mg + O_2 \rightarrow 2MgO$	Shows the number of atoms and molecules in the reaction, these need to be balanced.

Method	Description	Example
Filtration	<i>Separating an insoluble solid from a liquid</i>	To get sand from a mixture of sand, salt and water.
Crystallisation	<i>To separate a solid from a solution</i>	To obtain pure crystals of sodium chloride from salt water.
Simple distillation	<i>To separate a solvent from a solution</i>	To get pure water from salt water.
Fractional distillation	<i>Separating a mixture of liquids each with different boiling points</i>	To separate the different compounds in crude oil.
Chromatography	<i>Separating substances that move at different rates through a medium</i>	To separate out the dyes in food colouring.

Relative atomic mass

Isotopes	<i>Atoms of the same element with the same number of protons and different numbers of neutrons</i>	<p>^{35}Cl (75%) and ^{37}Cl (25%)</p> <p>Relative abundance = $(\% \text{ isotope 1} \times \text{mass isotope 1}) + (\% \text{ isotope 2} \times \text{mass isotope 2}) \div 100$ e.g. $(25 \times 37) + (75 \times 35) \div 100 = 35.5$</p>
-----------------	--	---

Alkali metals: 1, 2
 Halogens: 3, 4, 5, 6, 7
 Noble gases: 0

H	Transition metals										He						
Li	Be											B	C	N	O	F	Ne
Na	Mg											Al	Si	P	S	Cl	Ar
K	Ca	Sc	Ti	V	Cr	Mn	Fe	Co	Ni	Cu	Zn	Ga	Ge	As	Se	Br	Kr
Rb	Sr	Y	Zr	Nb	Mo	Tc	Ru	Rh	Pd	Ag	Cd	In	Sn	Sb	Te	I	Xe
Cs	Ba	La	Hf	Ta	W	Re	Os	Ir	Pt	Au	Hg	Tl	Pb	Bi	Po	At	Rn
Fr	Ra	Ac	Rf	Db	Sg	Bh	Hs	Mt	?	?	?						

Elements arranged in order of atomic number

Elements with similar properties are in columns called groups

Elements in the same group have the same number of outer shell electrons and elements in the same period (row) have the same number of electron shells.

The Periodic table

Development of the Periodic table

Before discovery of protons, neutrons and electrons	Elements arranged in order of atomic weight	Early periodic tables were incomplete, some elements were placed in inappropriate groups if the strict order atomic weights was followed.
Mendeleev	Left gaps for elements that hadn't been discovered yet	Elements with properties predicted by Mendeleev were discovered and filled in the gaps. Knowledge of isotopes explained why order based on atomic weights was not always correct.

Metals to the left of this line, non metals to the right

Metals	To the left of the Periodic table	Form positive ions. Conductors, high melting and boiling points, ductile, malleable.
Non metals	To the right of the Periodic table	Form negative ions. Insulators, low melting and boiling points.

Metals and non metals

Group 7

AQA GCSE Atomic structure and periodic table part 2

Group 0

Group 1

Alkali metals	Very reactive with oxygen, water and chlorine	Only have one electron in their outer shell. Form +1 ions.
	Reactivity increases down the group	Negative outer electron is further away from the positive nucleus so is more easily lost.

Halogens	Consist of molecules made of a pair of atoms	Have seven electrons in their outer shell. Form -1 ions.	
	Melting and boiling points increase down the group (gas → liquid → solid)	Increasing atomic mass number.	
	Reactivity decreases down the group	Increasing proton number means an electron is more easily gained	
With metals	Forms a metal halide	Metal + halogen → metal halide e.g. Sodium + chlorine → sodium chloride	e.g. NaCl metal atom loses outer shell electrons and halogen gains an outer shell electron
With hydrogen	Forms a hydrogen halide	Hydrogen + halogen → hydrogen halide e.g. Hydrogen + bromine → hydrogen bromide	e.g. Cl ₂ + H ₂ → 2HCl
With aqueous solution of a halide salt	A more reactive halogen will displace the less reactive halogen from the salt	Chlorine + potassium bromide → potassium chloride + bromine	e.g. Cl ₂ + 2KBr → 2KCl + Br ₂

Noble gases	Unreactive, do not form molecules	This is due to having full outer shells of electrons.
	Boiling points increase down the group	Increasing atomic number.

With oxygen	Forms a metal oxide	Metal + oxygen → metal oxide	e.g. 4Na + O ₂ → 2Na ₂ O
With water	Forms a metal hydroxide and hydrogen	Metal + water → metal hydroxide + hydrogen	e.g. 2Na + 2H ₂ O → 2NaOH + H ₂
With chlorine	Forms a metal chloride	Metal + chlorine → metal chloride	e.g. 2Na + Cl ₂ → 2NaCl

Ionic	<i>Particles are oppositely charged ions</i>	Occurs in compounds formed from metals combined with non metals.
Covalent	<i>Particles are atoms that share pairs of electrons</i>	Occurs in most non metallic elements and in compounds of non metals.
Metallic	<i>Particles are atoms which share delocalised electrons</i>	Occurs in metallic elements and alloys.

Solid, liquid, gas

Melting and freezing happen at melting point, boiling and condensing happen at boiling point.

The amount of energy needed for a state change depends on the strength of forces between particles in the substance.

<i>s</i>	solid
<i>l</i>	liquid
<i>g</i>	gas

Chemical bonds

The three states of matter

<i>Good conductors of electricity</i>	Delocalised electrons carry electrical charge through the metal.
<i>Good conductors of thermal energy</i>	Energy is transferred by the delocalised electrons.

<i>High melting and boiling points</i>	This is due to the strong metallic bonds.
<i>Pure metals can be bent and shaped</i>	Atoms are arranged in layers that can slide over each other.

<i>High melting and boiling points</i>	Large amounts of energy needed to break the bonds.
<i>Do not conduct electricity when solid</i>	Ions are held in a fixed position in the lattice and cannot move.
<i>Do conduct electricity when molten or dissolved</i>	Lattice breaks apart and the ions are free to move.

Properties of ionic compounds

AQA BONDING, STRUCTURE AND THE PROPERTIES OF MATTER 1

Metals as conductors

Properties of metals and alloys

Alloys	<i>Mixture of two or more elements at least one of which is a metal</i>	Harder than pure metals because atoms of different sizes disrupt the layers so they cannot slide over each other.
---------------	---	---

Ionic bonding

Electrons are transferred so that all atoms have a noble gas configuration (full outer shells).	<i>Metal atoms lose electrons and become positively charged ions</i>	Group 1 metals form +1 ions Group 2 metals form +2 ions
	<i>Non metals atoms gain electrons to become negatively charged ions</i>	Group 6 non metals form -2 ions Group 7 non metals form -1 ions

Pure metal **Alloy**

Metallic bonding

Dot and cross diagram

Giant structure

Ionic compounds

Structure

- Held together by strong electrostatic forces of attraction between oppositely charged ions*
- Forces act in all directions in the lattice*


Giant structure of atoms arranged in a regular pattern

Electrons in the outer shell of metal atoms are delocalised and free to move through the whole structure. This sharing of electrons leads to strong metallic bonds.

Very large molecules	<i>Solids at room temperature</i>	Atoms are linked by strong covalent bonds.	
----------------------	-----------------------------------	--	--

<i>Each carbon atom is bonded to four others</i>		Very hard.	Rigid structure.
		Very high melting point.	Strong covalent bonds.
		Does not conduct electricity.	No delocalised electrons.

Usually gases or liquids	<i>Covalent bonds in the molecule are strong but forces between molecules (intermolecular) are weak</i>	Low melting and boiling points.	Due to having weak intermolecular forces that easily broken.
		Do not conduct electricity.	Due to them molecules not having an overall electrical charge.
		Larger molecules have higher melting and boiling points.	Intermolecular forces increase with the size of the molecules.



Atoms share pairs of electrons	<i>Can be small molecules e.g. ammonia</i>	<p>Dot and cross : + Show which atom the electrons in the bonds come from - All electrons are identical</p>
		<p>2D with bonds: + Show which atoms are bonded together - It shows the H-C-H bond incorrectly at 90°</p>
		<p>3D ball and stick model: + Attempts to show the H-C-H bond angle is 109.5°</p>
<i>Can be giant covalent structures e.g. polymers</i>		

Graphene		Excellent conductor.	Contains delocalised electrons.
		Very strong.	Contains strong covalent bonds.

Single layer of graphite one atom thick

Graphene and fullerenes

Diamond, graphite, silicon dioxide	<i>Very high melting points</i>	Lots of energy needed to break strong, covalent bonds.
------------------------------------	---------------------------------	--

Fullerenes		Buckminsterfullerene, C ₆₀	Hexagonal rings of carbon atoms with hollow shapes. Can also have rings of five (pentagonal) or seven (heptagonal) carbon atoms.
		First fullerene to be discovered.	

Carbon nanotubes		Very thin and long cylindrical fullerenes	Used in electronics industry.
		Very conductive.	Reinforcing composite materials.
		High tensile strength.	Catalysts and lubricants.
		Large surface area to volume ratio.	

<i>Each carbon atom is bonded to three others forming layers of hexagonal rings with no covalent bonds between the layers</i>		Slippery.	Layers can slide over each other.
		Very high melting point.	Strong covalent bonds.
		Does conduct electricity.	Delocalised electrons between layers.

Graphite

M_r	<i>The sum of the relative atomic masses of the atoms in the numbers shown in the formula</i>	The sum of the M_r of the reactants in the quantities shown equals the sum of the M_r of the products in the quantities shown.	$2Mg + O_2 \rightarrow 2MgO$ $48g + 32g = 80g$ $80g = 80g$
-------	---	--	--

Mass appears to increase during a reaction	<i>One of the reactants is a gas</i>	Magnesium + oxygen \rightarrow magnesium oxide
Mass appears to decrease during a reaction	<i>One of the products is a gas and has escaped</i>	Calcium carbonate \rightarrow carbon dioxide + calcium oxide

Mass changes when a reactant or product is a gas

Relative formula mass (M_r)

Conservation of mass	<i>No atoms are lost or made during a chemical reaction</i>	Mass of the products equals the mass of the reactants.
----------------------	---	--

AQA GCSE QUANTITATIVE CHEMISTRY

Balanced symbol equations	<i>Represent chemical reactions and have the same number of atoms of each element on both sides of the equation</i>	$H_2 + Cl_2 \rightarrow 2HCl$ Subscript \swarrow \nwarrow Normal script Subscript numbers show the number of atoms of the element to its left. Normal script numbers show the number of molecules.
---------------------------	---	---

Conservation of mass and balanced symbol equations

Chemical measurements

Whenever a measurement is taken, there is always some uncertainty about the result obtained	<i>Can determine whether the mean value falls within the range of uncertainty of the result</i>	1. Calculate the mean 2. Calculate the range of the results 3. Estimate of uncertainty in mean would be half the range
---	---	--

Concentration of solutions	Example: 1. Mean value is 46.5s 2. Range of results is 44s to 49s = 5s 3. Time taken was $46.5s \pm 2.5s$
----------------------------	--

Measured in mass per given volume of solution (g/dm^3)	<i>Conc. = $\frac{mass (g)}{volume (dm^3)}$</i>	HT only Greater mass = higher concentration. Greater volume = lower concentration.
--	--	--

Acids	<i>Acids produce hydrogen ions (H⁺) in aqueous solutions.</i>
Alkalis	<i>Aqueous solutions of alkalis contain hydroxide ions (OH⁻).</i>

You can use universal indicator or a pH probe to measure the acidity or alkalinity of a solution against the pH scale.

In neutralisation reactions, hydrogen ions react with hydroxide ions to produce water:
 $H^+ + OH^- \rightarrow H_2O$

Reactions with acids	<i>metal + acid → metal salt + hydrogen</i>	magnesium + hydrochloric acid → magnesium chloride + hydrogen zinc + sulfuric acid → zinc sulfate + hydrogen
-----------------------------	---	---

Acids react with some metals to produce salts and hydrogen.

Extraction using carbon	
<i>Metals less reactive than carbon can be extracted from their oxides by reduction.</i>	For example: zinc oxide + carbon → zinc + carbon dioxide

Reactions of acids and metals

Extraction of metals and reduction

Unreactive metals, such as gold, are found in the Earth as the metal itself. They can be mined from the ground.

Reactions of acids

AQA Chemical Changes 1

Reactivity of metals

The reactivity series

	Reactions with water	Reactions with acid
Group 1 metals	<i>Reactions get more vigorous as you go down the group</i>	<i>Reactions get more vigorous as you go down the group</i>
Group 2 metals	<i>Do not react with water</i>	<i>Observable reactions include fizzing and temperature increases</i>
Zinc, iron and copper	<i>Do not react with water</i>	<i>Zinc and iron react slowly with acid. Copper does not react with acid.</i>

The pH scale and neutralisation

Soluble salts

Neutralisation of acids and salt production

Soluble salts	<i>Soluble salts can be made from reacting acids with solid insoluble substances (e.g. metals, metal oxides, hydroxides and carbonates).</i>
Production of soluble salts	<i>Add the solid to the acid until no more dissolves. Filter off excess solid and then crystallise to produce solid salts.</i>


Acid name	Salt name
<i>Hydrochloric acid</i>	Chloride
<i>Sulfuric acid</i>	Sulfate
<i>Nitric acid</i>	Nitrate

sodium hydroxide + hydrochloric acid → sodium chloride + water
calcium carbonate + sulfuric acid → calcium sulfate, + carbon dioxide + water

Neutralisation	<i>Acids can be neutralised by alkalis and bases</i>	An alkali is a soluble base e.g. metal hydroxide. A base is a substance that neutralises an acid e.g. a soluble metal hydroxide or a metal oxide.
-----------------------	--	--

Metal oxides

Metals form positive ions when they react	<i>The reactivity of a metal is related to its tendency to form positive ions</i>	The reactivity series arranges metals in order of their reactivity (their tendency to form positive ions).
Carbon and hydrogen	<i>Carbon and hydrogen are non-metals but are included in the reactivity series</i>	These two non-metals are included in the reactivity series as they can be used to extract some metals from their ores, depending on their reactivity.
Displacement	<i>A more reactive metal can displace a less reactive metal from a compound.</i>	Silver nitrate + Sodium chloride → Sodium nitrate + Silver chloride



Metals and oxygen	<i>Metals react with oxygen to form metal oxides</i>	magnesium + oxygen → magnesium oxide $2Mg + O_2 \rightarrow 2MgO$
Reduction	<i>This is when oxygen is removed from a compound during a reaction</i>	e.g. metal oxides reacting with hydrogen, extracting low reactivity metals
Oxidation	<i>This is when oxygen is gained by a compound during a reaction</i>	e.g. metals reacting with oxygen, rusting of iron

The ions discharged when an aqueous solution is electrolysed using inert electrodes depend on the relative reactivity of the elements involved.

At the negative electrode	Metal will be produced on the electrode if it is less reactive than hydrogen. Hydrogen will be produced if the metal is more reactive than hydrogen.
At the positive electrode	Oxygen is formed at positive electrode. If you have a halide ion (Cl ⁻ , I ⁻ , Br ⁻) then you will get chlorine, bromine or iodine formed at that electrode.

Electrolysis of aqueous solutions

Process of electrolysis	<i>Splitting up using electricity</i>	When an ionic compound is melted or dissolved in water, the ions are free to move. These are then able to conduct electricity and are called electrolytes. Passing an electric current through electrolytes causes the ions to move to the electrodes.
Electrode	<i>Anode Cathode</i>	The positive electrode is called the anode. The negative electrode is called the cathode.
Where do the ions go?	<i>Cations Anions</i>	Cations are positive ions and they move to the negative cathode. Anions are negative ions and they move to the positive anode.

Extracting metals using electrolysis	<i>Metals can be extracted from molten compounds using electrolysis.</i>
	<i>This process is used when the metal is too reactive to be extracted by reduction with carbon.</i>
	<i>The process is expensive due to large amounts of energy needed to produce the electrical current. Example: aluminium is extracted in this way.</i>

Electrolysis

AQA Chemical Changes 2

AQA GCSE Energy changes

Reaction profiles

Activation energy	<i>Chemical reactions only happen when particles collide with sufficient energy</i>	The minimum amount of energy that colliding particles must have in order to react is called the activation energy.
--------------------------	---	--

Types of reaction

Endothermic	<i>Energy is taken in from the surroundings so the temperature of the surroundings decreases</i>	<ul style="list-style-type: none"> Thermal decomposition Sports injury packs
Exothermic	<i>Energy is transferred to the surroundings so the temperature of the surroundings increases</i>	<ul style="list-style-type: none"> Combustion Hand warmers Neutralisation

Endothermic		Products are at a higher energy level than the reactants. As the reactants form products, energy is transferred from the surroundings to the reaction mixture. The temperature of the surroundings decreases because energy is taken in during the reaction.
Exothermic		Products are at a lower energy level than the reactants. When the reactants form products, energy is transferred to the surroundings. The temperature of the surroundings increases because energy is released during the reaction.

Rate of chemical reaction	<i>This can be calculated by measuring the quantity of reactant used or product formed in a given time.</i>	Rate = $\frac{\text{quantity of reactant used}}{\text{time taken}}$ Rate = $\frac{\text{quantity of product formed}}{\text{time taken}}$
---------------------------	---	---

Factors affecting the rate of reaction	
Temperature	<i>The higher the temperature, the quicker the rate of reaction.</i>
Concentration	<i>The higher the concentration, the quicker the rate of reaction.</i>
Surface area	<i>The larger the surface area of a reactant solid, the quicker the rate of reaction.</i>
Pressure (of gases)	<i>When gases react, the higher the pressure upon them, the quicker the rate of reaction.</i>


Quantity	Unit
<i>Mass</i>	Grams (g)
<i>Volume</i>	cm ³
<i>Rate of reaction</i>	Grams per cm ³ (g/cm ³) HT: moles per second (mol/s)

Calculating rates of reactions

Rate of reaction

Factors affecting rates


Collision theory and activation energy



**AQA GCSE
The rate and extent of chemical change**

Reversible reactions and dynamic equilibrium

Collision theory	<i>Chemical reactions can only occur when reacting particles collide with each other with sufficient energy.</i>	Increasing the temperature increases the frequency of collisions and makes the collisions more energetic, therefore increasing the rate of reaction.
Activation energy	<i>This is the minimum amount of energy colliding particles in a reaction need in order to react.</i>	Increasing the concentration, pressure (gases) and surface area (solids) of reactions increases the frequency of collisions, therefore increasing the rate of reaction.



If a catalyst is used in a reaction, it is not shown in the word equation.

Catalyst	A catalyst changes the rate of a chemical reaction but is not used in the reaction.
Enzymes	These are biological catalysts.
How do they work?	Catalysts provide a different reaction pathway where reactants do not require as much energy to react when they collide.

Catalysts

Reversible reactions


Reversible reactions	In some chemical reactions, the products can react again to re-form the reactants.
Representing reversible reactions	$A + B \rightleftharpoons C + D$
The direction	The direction of reversible reactions can be changed by changing conditions: $A + B \xrightleftharpoons[\text{cool}]{\text{heat}} C + D$

Equilibrium

Equilibrium in reversible reactions
When a reversible reaction occurs in apparatus which prevents the escape of reactants and products, equilibrium is reached when the forward and reverse reactions occur exactly at the same rate.


Energy changes and reversible reactions

If one direction of a reversible reaction is exothermic, the opposite direction is endothermic. The same amount of energy is transferred in each case.



Crude oil, hydrocarbons and alkanes

Display formula for first four alkanes



Fractions	<i>The hydrocarbons in crude oil can be split into fractions</i>	Each fraction contains molecules with a similar number of carbon atoms in them. The process used to do this is called fractional distillation.
Using fractions	<i>Fractions can be processed to produce fuels and feedstock for petrochemical industry</i>	We depend on many of these fuels; petrol, diesel and kerosene. Many useful materials are made by the petrochemical industry; solvents, lubricants and polymers.


Carbon compounds as fuels and feedstock

AQA GCSE Organic chemistry 1

Carbon compounds as fuels and feedstock

Fractional distillation and petrochemicals

Hydrocarbon chains	In oil	Hydrocarbon chains in crude oil come in lots of different lengths.
	Boiling points	The boiling point of the chain depends on its length. During fractional distillation, they boil and separate at different temperatures due to this.



Crude oil	<i>A finite resource</i>	Consisting mainly of plankton that was buried in the mud, crude oil is the remains of ancient biomass.
Hydrocarbons	<i>These make up the majority of the compounds in crude oil</i>	Most of these hydrocarbons are called alkanes.
General formula for alkanes	<i>C_nH_{2n+2}</i>	For example: C ₂ H ₆ C ₆ H ₁₄



Alkanes to alkenes	<i>Long chain alkanes are cracked into short chain alkenes.</i>
Alkenes	<i>Alkenes are hydrocarbons with a double bond (some are formed during the cracking process).</i>
Properties of alkenes	<i>Alkenes are more reactive than alkanes and react with bromine water. Bromine water changes from orange to colourless in the presence of alkenes.</i>

Cracking and alkenes

Properties of hydrocarbons


Combustion	During the complete combustion of hydrocarbons, the carbon and hydrogen in the fuels are oxidised, releasing carbon dioxide, water and energy.
-------------------	--

Cracking	<i>The breaking down of long chain hydrocarbons into smaller chains</i>	The smaller chains are more useful. Cracking can be done by various methods including catalytic cracking and steam cracking.
Catalytic cracking	<i>The heavy fraction is heated until vaporised</i>	After vaporisation, the vapour is passed over a hot catalyst forming smaller, more useful hydrocarbons.
Steam cracking	<i>The heavy fraction is heated until vaporised</i>	After vaporisation, the vapour is mixed with steam and heated to a very high temperature forming smaller, more useful hydrocarbons.



Alkenes and uses as polymers	<i>Used to produce polymers. They are also used as the starting materials of many other chemicals, such as alcohol, plastics and detergents.</i>	Boiling point (temperature at which liquid boils)	<i>As the hydrocarbon chain length increases, boiling point increases.</i>
Why do we crack long chains?	<i>Without cracking, many of the long hydrocarbons would be wasted as there is not much demand for these as for the shorter chains.</i>	Viscosity (how easily it flows)	<i>As the hydrocarbon chain length increases, viscosity increases.</i>
		Flammability (how easily it burns)	<i>As the hydrocarbon chain length increases, flammability decreases.</i>

Pure substances	<i>A pure substance is a single element or compound, not mixed with any other substance.</i>	Pure substances melt and boil at specific temperatures. Heating graphs can be used to distinguish pure substances from impure.
------------------------	--	--



Pure substances

Purity, formulations and chromatography


Chromatography

Formulations

AQA Chemical analysis


Identification of common gases

Formulation	<i>A formulation is a mixture that has been designed as a useful product.</i>
How are formulations made?	<i>By mixing chemicals that have a particular purpose in careful quantities.</i>
Examples of formulations.	<i>Fuels, cleaning agents, paints, medicines and fertilisers.</i>



Chromatography	<i>Can be used to separate mixtures and help identify substances.</i>	Involves a mobile phase (e.g. water or ethanol) and a stationary phase (e.g. chromatography paper).
R_f Values	<i>The ratio of the distance moved by a compound to the distance moved by solvent.</i>	$R_f = \frac{\text{distance moved by substance}}{\text{distance moved by solvent}}$
Pure substances	<i>The compounds in a mixture separate into different spots.</i>	This depends on the solvent used. A pure substance will produce a single spot in all solvents whereas an impure substance will produce multiple spots.

Gas	Test	Positive result
Hydrogen	<i>Burning splint</i>	'Pop' sound.
Oxygen	<i>Glowing splint</i>	Re-lights the splint.
Chlorine	<i>Litmus paper (damp)</i>	Bleaches the paper white.
Carbon dioxide	<i>Limewater</i>	Goes cloudy (as a solid calcium carbonate forms).



Gas	Percentage
Nitrogen	~80%
Oxygen	~20%
Argon	0.93%
Carbon dioxide	0.04%

Proportions of gases in the atmosphere

Algae and plants	<i>These produced the oxygen that is now in the atmosphere, through photosynthesis.</i>	carbon dioxide + water → glucose + oxygen $6\text{CO}_2 + 6\text{H}_2\text{O} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6\text{O}_2$
Oxygen in the atmosphere	<i>First produced by algae 2.7 billion years ago.</i>	Over the next billion years plants evolved to gradually produce more oxygen. This gradually increased to a level that enabled animals to evolve.

The Earth's early atmosphere

How oxygen increased
How carbon dioxide decreased

Composition and evolution of the atmosphere

AQA GCSE Chemistry of the atmosphere

Common atmospheric pollutants

CO₂ and methane as greenhouse gases

Carbon footprints
The total amount of greenhouse gases emitted over the full life cycle of a product/event. This can be reduced by reducing emissions of carbon dioxide and methane.

Global climate change

Greenhouse gases

Carbon dioxide, water vapour and methane	<i>Examples of greenhouse gases that maintain temperatures on Earth in order to support life</i>
The greenhouse effect	<i>Radiation from the Sun enters the Earth's atmosphere and reflects off of the Earth. Some of this radiation is re-radiated back by the atmosphere to the Earth, warming up the global temperature.</i>

Human activities and greenhouse gases

Carbon dioxide	<i>Human activities that increase carbon dioxide levels include burning fossil fuels and deforestation.</i>
Methane	<i>Human activities that increase methane levels include raising livestock (for food) and using landfills (the decay of organic matter released methane).</i>
Climate change	<i>There is evidence to suggest that human activities will cause the Earth's atmospheric temperature to increase and cause climate change.</i>

Effects of climate change
Rising sea levels
Extreme weather events such as severe storms
Change in amount and distribution of rainfall
Changes to distribution of wildlife species with some becoming extinct

Properties and effects of atmospheric pollutants	
Carbon monoxide	<i>Toxic, colourless and odourless gas. Not easily detected, can kill.</i>
Sulfur dioxide and oxides of nitrogen	<i>Cause respiratory problems in humans and acid rain which affects the environment.</i>
Particulates	<i>Cause global dimming and health problems in humans.</i>

Atmospheric pollutants from fuels

Combustion of fuels	<i>Source of atmospheric pollutants. Most fuels may also contain some sulfur.</i>
Gases from burning fuels	<i>Carbon dioxide, water vapour, carbon monoxide, sulfur dioxide and oxides of nitrogen.</i>
Particulates	<i>Solid particles and unburned hydrocarbons released when burning fuels.</i>

Volcano activity 1 st Billion years	<i>Billions of years ago there was intense volcanic activity</i>	This released gases (mainly CO ₂) that formed to early atmosphere and water vapour that condensed to form the oceans.
Other gases	<i>Released from volcanic eruptions</i>	Nitrogen was also released, gradually building up in the atmosphere. Small proportions of ammonia and methane also produced.
Reducing carbon dioxide in the atmosphere	<i>When the oceans formed, carbon dioxide dissolved into it</i>	This formed carbonate precipitates, forming sediments. This reduced the levels of carbon dioxide in the atmosphere.

Sterilising agents include chlorine, ozone and UV light.

Potable water	<i>Water of an appropriate quality is essential for life</i>	Human drinking water should have low levels of dissolved salts and microbes. This is called potable water.
UK water	<i>Rain provides water with low levels of dissolved substances</i>	This water collects in the ground/lakes/streams. To make potable water an appropriate source is chosen, which is then passed through filter beds and then sterilised.
Desalination	<i>Needs to occur is fresh water is limited and salty/sea water is needed for drinking</i>	This can be achieved by distillation or by using large membranes e.g. reverse osmosis. These processes require large amounts of energy.

Earth's resources	<i>Used to provide warmth, shelter, food and transport for humans</i>	Natural resources and resources from agriculture provide: timber, food, clothing and fuels.
		Finite resources from the Earth, oceans and atmosphere are processed to provide energy and materials.
Chemistry and resources	<i>Research and techniques improve agricultural and industrial processes</i>	These improvements provide new products and improve sustainability.
Plastics	<i>Normally made using ethene from crude oil</i>	However, the raw material ethene can also be obtained from ethanol, which can be produced during fermentation. Industries are now starting to use a renewable crop for this process.

Using the Earth's resources and sustainable development

Potable water

Using the Earth's resources and obtaining potable water

AQA GCSE Using resources 1

Life cycle assessment and recycling

Ways of reducing the use of resources

Waste water treatment

Waste water	<i>Produced from urban lifestyles and industrial processes</i>	These require treatment before used in the environment. Sewage needs the organic matter and harmful microbes removed.
Sewage treatment	<i>Includes many stages</i>	<ul style="list-style-type: none"> - Screening and grit removal - Sedimentation to produce sludge and effluent (liquid waste or sewage). - Anaerobic digestion of sludge - Aerobic biological treatment of effluent.

LCAS	<i>Life cycle assessments are carried out to assess the environmental impact of products</i>	They are assessed at these stages: <ul style="list-style-type: none"> - Extraction and processing raw materials - Manufacturing and packaging - Use and operation during lifetime - Disposal
Values	<i>Allocating numerical values to pollutant effects is difficult</i>	Value judgments are allocated to the effects of pollutants so LCA is not a purely objective process.

Life cycle assessment

Reduce, reuse and recycle	<i>This strategy reduces the use of limited resources</i>	This, therefore, reduces energy sources being used, reduces waste (landfill) and reduces environmental impacts.
Limited raw materials	<i>Used for metals, glass, building materials, plastics and clay ceramics</i>	Most of the energy required for these processes comes from limited resources. Obtaining raw materials from the Earth by quarrying and mining causes environmental impacts.
Reusing and recycling	<i>Metals can be recycled by melting and recasting/reforming</i>	Glass bottles can be reused. They are crushed and melted to make different glass products. Products that cannot be reused are recycled.

Mechanical	<i>Force acts upon an object</i>
Electrical	<i>Electric current flow</i>
Heat	<i>Temperature difference between objects</i>
Radiation	<i>Electromagnetic waves or sound</i>


Energy pathways

Change in thermal energy = mass X specific heat capacity X temperature change $\Delta E = m \times c \times \Delta \theta$

Specific Heat Capacity
Energy needed to raise 1kg of substance by 1°C
Depends on: mass of substance, what the substance is and energy put into the system.

Efficiency = $\frac{\text{Useful power output}}{\text{Total power input}}$

Efficiency = $\frac{\text{Useful output energy transfer}}{\text{Total input energy transfer}}$



Kinetic energy	<i>Energy stored by a moving object</i>	$\frac{1}{2} \times \text{mass} \times (\text{speed})^2$ $\frac{1}{2} mv^2$
Elastic Potential energy	<i>Energy stored in a stretched spring, elastic band</i>	$\frac{1}{2} \times \text{spring constant} \times (\text{extension})^2$ $\frac{1}{2} ke^2$ (Assuming the limit of proportionality has not been exceeded)
Gravitational Potential energy	<i>Energy gained by an object raised above the ground</i>	Mass X gravitational field strength X height mgh

System	<i>An object or group of objects that interact together</i>	EG: Kettle boiling water.
Energy stores	<i>Kinetic, chemical, internal (thermal), gravitational potential, elastic potential, magnetic, electrostatic, nuclear</i>	Energy is gained or lost from the object or device.
Ways to transfer energy	<i>Light, sound, electricity, thermal, kinetic are ways to transfer from one store to another store of energy.</i>	EG: electrical energy transfers chemical energy into thermal energy to heat water up.
Unit	<i>Joules (J)</i>	

Work	<i>Doing work transfers energy from one store to another</i>	By applying a force to move an object the energy store is changed.	Work done = Force X distance moved $W = Fs$
Power	<i>The rate of energy transfer</i>	1 Joule of energy per second = 1 watt of power	Power = energy transfer ÷ time $P = E \div t$ Power = work done ÷ time, $P = W \div t$

	<i>Units</i>
Specific Heat Capacity	<i>Joules per Kilogram degree Celsius (J/Kg°C)</i>
Temperature change	<i>Degrees Celsius (°C)</i>
Work done	<i>Joules (J)</i>
Force	<i>Newton (N)</i>
Distance moved	<i>Metre (m)</i>
Power	<i>Watts (W)</i>
Time	<i>Seconds (s)</i>

Useful energy	<i>Energy transferred and used</i>
Wasted energy	<i>Dissipated energy, stored less usefully</i>


Prefix	<i>Multiple</i>	Standard form
Kilo	<i>1000</i>	10^3
Mega	<i>1000 000</i>	10^6
Giga	<i>100 000 000</i>	10^9

Energy stores and changes

AQA ENERGY – part 1

Energy Conservation and Dissipation

Closed system	<i>No change in total energy in system</i>
Open system	<i>Energy can dissipate</i>



Dissipate
To scatter in all directions or to use wastefully
When energy is 'wasted', it dissipates into the surroundings as internal (thermal) energy.



Ways to reduce 'wasted' energy
Energy transferred usefully
Insulation, streamline design, lubrication of moving parts.

Principle of conservation of energy
The amount of energy always stays the same.
Energy cannot be created or destroyed, only changed from one store to another.

	<i>Units</i>
Energy (KE, EPE, GPE, thermal)	<i>Joules (J)</i>
Velocity	<i>Metres per second (m/s)</i>
Spring constant	<i>Newton per metre (N/m)</i>
Extension	<i>Metres (m)</i>
Mass	<i>Kilogram (Kg)</i>
Gravitational field strength	<i>Newton per kilogram (N/Kg)</i>
Height	<i>Metres (m)</i>

Using renewable energy will need to increase to meet demand.

Renewable energy makes up about 20% of energy consumption.

Fossil fuel reserves are running out.

Energy demand is increasing as population increases.

Non-renewable energy resource	These will run out. It is a finite reserve. It cannot be replenished.	e.g. Fossil fuels (coal, oil and gas) and nuclear fuels.
Renewable energy resource	These will never run out. It is an infinite reserve. It can be replenished.	e.g. Solar, Tides, Waves, Wind, Geothermal, Biomass, Hydroelectric


Using fuels

Energy resources


Global Energy Resources

AQA ENERGY – part 2


National Grid



Power station – NB: You need to understand the principle behind generating electricity. An energy resource is burnt to make steam to drive a turbine which drives the generator.




Energy resource	How it works	Uses	Positive	Negative
Fossil Fuels (coal, oil and gas)	Burnt to release thermal energy used to turn water into steam to turn turbines	Generating electricity, heating and transport	Provides most of the UK energy. Large reserves. Cheap to extract. Used in transport, heating and making electricity. Easy to transport.	Non-renewable. Burning coal and oil releases sulfur dioxide. When mixed with rain makes acid rain. Acid rain damages building and kills plants. Burning fossil fuels releases carbon dioxide which contributes to global warming. Serious environmental damage if oil spilt.
Nuclear	Nuclear fission process	Generating electricity	No greenhouse gases produced. Lots of energy produced from small amounts of fuel.	Non-renewable. Dangers of radioactive materials being released into air or water. Nuclear sites need high levels of security. Start up costs and decommission costs very expensive. Toxic waste needs careful storing.
Biofuel	Plant matter burnt to release thermal energy	Transport and generating electricity	Renewable. As plants grow, they remove carbon dioxide. They are 'carbon neutral'.	Large areas of land needed to grow fuel crops. Habitats destroyed and food not grown. Emits carbon dioxide when burnt thus adding to greenhouse gases and global warming.
Tides	Every day tides rise and fall, so generation of electricity can be predicted	Generating electricity	Renewable. Predictable due to consistency of tides. No greenhouse gases produced.	Expensive to set up. A dam like structure is built across an estuary, altering habitats and causing problems for ships and boats.
Waves	Up and down motion turns turbines	Generating electricity	Renewable. No waste products.	Can be unreliable depends on wave output as large waves can stop the pistons working.
Hydroelectric	Falling water spins a turbine	Generating electricity	Renewable. No waste products.	Habitats destroyed when dam is built.
Wind	Movement causes turbine to spin which turns a generator	Generating electricity	Renewable. No waste products.	Unreliable – wind varies. Visual and noise pollution. Dangerous to migrating birds.
Solar	Directly heats objects in solar panels or sunlight captured in photovoltaic cells	Generating electricity and some heating	Renewable. No waste products.	Making and installing solar panels expensive. Unreliable due to light intensity.
Geothermal	Hot rocks under the ground heats water to produce steam to turn turbine	Generating electricity and heating	Renewable. Clean. No greenhouse gases produced.	Limited to a small number of countries. Geothermal power stations can cause earthquake tremors.



Electrons carry current.
Electrons are free to move in metal.

Cell	Battery	Switch	Lamp	Ammeter	Volt meter	Diode	LED	LDR	Fuse	Resistor	Variable resistor	Thermistor
<i>Store of chemical energy</i>	<i>Two or more cells in series</i>	<i>Breaks circuit, turning current off</i>	<i>Lights when current flows</i>	<i>Measures current</i>	<i>Measures potential difference</i>	<i>Current flows one way</i>	<i>Emits light when current flows</i>	<i>Resistance low in bright light</i>	<i>Melts when current is too high</i>	<i>Affects the size of current flowing</i>	<i>Allows current to be varied</i>	<i>Resistance low at high temp</i>



Current	<i>Flow of electrical charge</i>	Ampere (A)
Potential difference (p.d.)	<i>How much electrical work is done by a cell</i>	Volts (V)
Charge	<i>Amount of electricity travelling in a circuit</i>	Coulombs (C)

Circuit symbols

Current and Charge
Current, potential difference and resistance

Series and parallel circuits

Series circuit	Current is the same in all components.	Total p.d. from battery is shared between all the components.	Total resistance is the sum of each component's resistance.
Parallel circuit	Total current is the sum of each component's current.	p.d. across all components is the same.	Total resistance is less than the resistance value of the smallest individual resistor.


Series	Parallel
<i>A circuit with one loop</i>	<i>A circuit with two or more loops</i>

Total p.d.
If cells are joined in series, add up individual cell values

Charge = Current X time $Q = I \times t$

Controlling current

Changing current
Change the p.d. of the cells
Add more components



$R = V \div I$

Resistance = Potential difference ÷ Current

AQA Electricity

Energy transfers

Work is done when charge flowing.

Power (W) = potential difference X current $R = V \times I$
 Power = (current)² X resistance $P = I^2 \times R$
 Energy transferred = Power X time $E = P \times t$

Domestic uses and safety

National Grid
Distributes electricity generated in power stations around UK

Step-up transformers	Step-down transformers
<i>Increase voltage, decrease current</i>	<i>Decrease voltage, increase current</i>
Increases efficiency, reduces heat loss.	Makes safer for houses.

Ammeter	<i>Set up in series with components</i>
Voltmeter	<i>Set up parallel to components</i>
Resistance (Ω)	<i>A measurement of how much current flow is reduced</i>
The higher the resistance, the more difficult it is for current to flow.	
Increasing resistance, reduces current.	
Increasing voltage, increases current.	

Thermistor	LDR	Alternating current	Direct current
<i>Resistance varies with temperature</i>	<i>Resistance varies with light intensity</i>	<i>p.d. switches direction many times a second, current switches direction</i>	<i>p.d. remains in one direction, current flows the same direction</i>
Resistance decreases as temperature increases.	Resistance decreases as light increases.	Generator.	Cell or battery.


Ohmic conductor or	<i>At a constant temperature, current is directly proportional to the p.d. across the resistor.</i>
Filament lamp	<i>As current increases, the resistance increases. The temperature increases as current flows.</i>
Diode	<i>Current flows when p.d. flows forward. Very high resistance in reverse.</i>


Current – Potential difference graphs

'Earthing' a safety device; Earth wire joins the metal case.

Mains supply
Frequency 50Hz, 230V

3 pin plug	<i>Live - Brown</i>	Carries p.d from mains supply.	p.d between live and earth = 230V
	<i>Neutral - Blue</i>	Completes the circuit.	p.d. = 0V
	<i>Earth - Green and Yellow stripes</i>	Only carries current if there is a fault.	p.d. = 0V





Pressure of a fixed volume of gas increases as temperature increases (temperature increases, speed increases, collisions occur more frequently and with more force so pressure increases).

Temperature of gas is linked to the average kinetic energy of the particles.

If kinetic energy increases so does the temperature of gas.

No kinetic energy is lost when gas particles collide with each other or the container.

Gas particles are in a constant state of random motion.

$$P = m \div V$$

$$\text{Density} = \text{mass} \div \text{volume.}$$

Density *Mass of a substance in a given volume*

Kinetic theory of gases

State	Particle arrangement	Properties
Solid	<i>Packed in a regular structure. Strong forces hold in place so cannot move.</i>	Difficult to change shape.
Liquid	<i>Close together, forces keep contact but can move about.</i>	Can change shape but difficult to compress.
Gas	<i>Separated by large distances. Weak forces so constantly randomly moving.</i>	Can expand to fill a space, easy to compress.

	Units
Density	<i>Kilograms per metre cubed (kg/m³)</i>
Mass	<i>Kilograms (kg)</i>
Volume	<i>Metres cubed (m³)</i>
Energy needed	<i>Joules (J)</i>
Specific latent heat	<i>Joule per kilogram (J/kg)</i>
Change in thermal energy	<i>Joules (J)</i>
Specific heat capacity	<i>Joule per kilogram degrees Celsius (J/kg°C)</i>
Temperature change	<i>Degrees Celsius (°C)</i>
Pressure	<i>Pascals (Pa)</i>

Particle model

AQA PARTICLE MODEL OF MATTER

Internal energy and energy transfers

Change of state

Specific Heat Capacity
Energy needed to raise 1kg of substance by 1°C
Depends on:
• Mass of substance
• What the substance is
• Energy put into the system.

Change in thermal energy = mass X specific heat capacity X temperature change.



$$\Delta E = m \times c \times \Delta \theta$$

Internal energy
Energy stored inside a system by particles
Internal energy is the total kinetic and potential energy of all the particles (atoms and molecules) in a system.
Heating changes the energy stored within a system
Heating causes a change in state. As particles separate, potential energy stored increases. Heating increases the temperature of a system. Particles move faster so kinetic energy of particles increases.

Specific Latent Heat	<i>Energy needed to change 1kg of a substance's state</i>
Specific Latent Heat of Fusion	<i>Energy needed to change 1kg of solid into 1 kg of liquid at the same temperature</i>
Specific Latent Heat of Vaporisation	<i>Energy needed to change 1kg of liquid into 1 kg of gas at the same temperature</i>

$$\text{Energy needed} = \text{mass} \times \text{specific latent heat.}$$

$$\Delta E = m \times L$$




Freezing	Liquid turns to a solid. Internal energy decreases.
Melting	Solid turns to a liquid. Internal energy increases.
Boiling / Evaporating	Liquid turns to a gas. Internal energy increases.
Condensation	Gas turns to a liquid. Internal energy decreases.
Sublimation	Solid turns directly into a gas. Internal energy increases.
Conservation of mass	When substances change state, mass is conserved.
Physical change	No new substance is made, process can be reversed.

Radius of an atom
 $1 \times 10^{-10} \text{m}$




Electrons gained
Negative ion

Electrons lost
Positive ion



Decay	Range in air	Ionising power	Penetration power
Alpha	Few cm	Very strong	Stopped by paper
Beta	Few m	Medium	Stopped by Aluminium
Gamma	Great distances	Weak	Stopped by thick lead



Atom	Same number of protons and electrons
Ion	Unequal number of electrons to protons
Mass number	Number of protons and neutrons
Atomic number	Number of protons

Particle	Charge	Size	Found
Neutron	None	1	In the nucleus
Proton	+	1	
Electron	-	Tiny	Orbits the nucleus


Atom structure

Isotope	${}^6_3\text{Li}$		${}^7_3\text{Li}$	
Different forms of an element with the same number of protons but different number of neutrons				

Discovery of the nucleus

Radioactive decay	Unstable atoms randomly emit radiation to become stable
Detecting	Use Geiger Muller tube
Unit	Becquerel
Ionisation	All radiation ionises

Decay	Emitted from nucleus	Changes in mass number and atomic number	
Alpha (α)	Helium nuclei (${}^4_2\text{He}$)	-4	-2
Beta (β)	Electron (${}^0_{-1}\text{e}$)	0	+1
Gamma (γ)	Electromagnetic wave	0	0
Neutron	Neutron	-1	0



Atoms and Isotopes

Atoms and Nuclear Radiation

AQA ATOMIC STRUCTURE

Democritus	Suggested idea of atoms as small spheres that cannot be cut.
J J Thomson (1897)	Discovered electrons– emitted from surface of hot metal. Showed electrons are negatively charged and that they are much less massive than atoms.
Thomson (1904)	Proposed 'plum pudding' model – atoms are a ball of positive charge with negative electrons embedded in it.
Geiger and Marsden (1909)	Directed beam of alpha particles (He^{2+}) at a thin sheet of gold foil. Found some travelled through, some were deflected, some bounced back.
Rutherford (1911)	Used above evidence to suggest alpha particles deflected due to electrostatic interaction between the very small charged nucleus, nucleus was massive. Proposed mass and positive charge contained in nucleus while electrons found outside the nucleus which cancel the positive charge exactly.
Bohr (1913)	Suggested modern model of atom – electrons in circular orbits around nucleus, electrons can change orbits by emitting or absorbing electromagnetic radiation. His research led to the idea of some particles within the nucleus having positive charge; these were named protons.
Chadwick (1932)	Discovered neutrons in nucleus – enabling other scientists to account for mass of atom.

Each Kg has a gravitational pull of 9.8N.

Unit	Newton (N)	1N
Kilo	Kilonewton (KN) = 1000	1X 10 ³
Mega	Meganewton (MN) = 1000,000	1 X 10 ⁶

Force	Push or pull	Stretch, squash, turn.
Contact force	Exerted between two objects when they touch	Friction, air resistance, tension.
Non-contact force	Exerted between two objects without touching	Gravity, electrostatic forces, magnetic forces.

Centre of mass **The weight of an object acts through a single point**

Gravitational field strength
Gravity exerted around an object.
Earth's gfs = 9.8N/kg

Weight = mass X gravitational field strength $W = m \times g$

Weight	Force acting upon an object due to gravity	Newton (N)
Mass	How much matter	Kilograms (Kg)


Gravity

Resultant force
The overall effect of all of the forces acting upon an object

Two forces acting in the same direction are added.
Two forces acting in the opposite direction are taken away.

Work done against frictional forces, temperature of object rises.

Object moves left with a force of 5N



Forces and their interactions

Contact and Resultant forces

If force is at right angles to direction of movement, NO work is done.

Scalar	A quantity that only has magnitude (size)	e.g. mass, time, speed, temperature, energy,
Vector	A quantity that only has magnitude and direction	e.g. force, velocity, momentum

Scalar and vector quantities

AQA FORCES – part 1

Work done and energy transfer

Work done
When work is done, energy is transferred
Work done = force X distance moved
 $W = F \times s$
1J of work is done when 1N of force moves an object through a distance of 1m, in the direction of the force.

An arrow can be used to show vectors

Length of arrow = magnitude of vector
Direction of arrow = direction of vector



Forces and elasticity

One force	The object changes speed or direction	Two balanced forces can stretch a object.
More than one force	The object changes shape	Two balanced forces can compress an object.
Elastic deformation	The object has been stretched but returns to its original length	Three balanced forces can bend an object.
Inelastic deformation	The object has been stretched but does not return to its original length	Limit of proportionality
Extension	The difference between stretched and unstretched lengths	Beyond this point the spring is permanently deformed

Velocity	Speed + direction	The speed of a car is 30m/s. A car moves forward with a velocity of 30m/s
Distance	How far	The table is 1m long
Displacement	Distance + direction	The beach is 1km due east of the town

Stretching a spring
Force = spring constant X extension, $F = k \times e$
 $EPE = \frac{1}{2} \times \text{spring constant} \times (\text{extension})^2$, $EPE = \frac{1}{2} ke^2$

Elastic Potential energy (EPE) Energy stored in a stretched spring

Force	Newton (N)
Spring constant	Newton per metre (N/m)
Extension	Metres (m)
EPE	Joules (J)

Area	Metres squares (m²)
Weight	Newton (N)
Mass	Kilograms (kg)
Gravitational field strength	Newton per kilogram (N/Kg)
Force	Newton (N)
Work done	Joules (J)
Distance	Metres (m)

Constant acceleration
 $(\text{final velocity})^2 - (\text{initial velocity})^2 = 2 \times \text{acceleration} \times \text{distance}$
 $v^2 - u^2 = 2 \times a \times s$

Velocity-time graph	<i>Shows speed of an object</i>	Falling objects	Falling objects accelerate due to gravity.	In no air resistance, objects accelerate at 9.8m/s^2	Air resistance slows falling objects down.
Accelerating	<i>Object getting faster</i>		Terminal velocity	<i>Weight of an object is balanced by resistive forces</i>	Object moves at a constant velocity. Resultant force = 0.
Decelerating	<i>Object slowing down</i>				
Acceleration = change in velocity ÷ time taken					
Acceleration	<i>Change in velocity</i>	Vector			


Velocity *The speed of an object with direction* Vector
 Speed of sound 330m/s

Speed = distance ÷ time
 $v = s \div t$

Speed	<i>How fast an object moves</i>	Scalar
Displacement	<i>Includes the distance and direction an object moves</i>	vector
Distance	<i>How far an object moves</i>	scalar

Distance-time graph *Shows how far an object moves along a straight line*
 Speed of object *Use the gradient of graph*

Forces, acceleration and Newton's Laws of motion



Speed or direction only changes if a resultant force acts on the object

Car on motorway	30m/s	Walking	1.5m/s
Train	60m/s	Running	3m/s
Jet plane	200m/s	Cycling	6m/s

Describing motion

Speed is rarely constant.

AQA FORCES – part 2
Observing and recording motion

Force = mass X acceleration
 $F = m \times a$

Acceleration is proportional to resultant force.
 Acceleration is inversely proportional to mass.

Newton's first Law	Balanced forces	When the resultant force on an still object = 0, the object is stationary. When the resultant force on a moving object = 0, the object is at a constant speed.
Newton's second Law	Unbalanced forces	When the resultant force is greater than 0, the object accelerates. It could speed up, slow down or change direction.
Newton's third Law	Equal and opposite forces	When two objects interact the forces exerted are equal and in an opposite direction.

Forces and braking

Speed affects both thinking and braking distances.
 Typical reaction time = 0.7s

Frictional forces decelerate a moving object and bring it to rest.

Thinking distance	<i>Distance travelled whilst the driver reacts</i>
Braking distance	<i>Distance travelled whilst the car is stopped by the brakes</i>
Stopping distance	<i>Total thinking and braking distances</i>

Factors affecting stopping distances	Drivers reaction times	Drinking alcohol, taking drugs, tired.
	Braking distances	Weather conditions, worn brakes or tyres, road surface, size of braking force.


Braking and kinetic energy
Work done by braking force, reduces kinetic energy
 Kinetic energy decreases, temperature of brakes increases due to frictional forces.

Speed / velocity	Metres per second (m/s)
Distance	Metres (m)
Time	Seconds (s)
Acceleration	Metres per second squared (m/s²)
Force	Newton (N)
Mass	Kilogram (Kg)

Relay

A device using a small current to control a larger current in another circuit

Solenoid is wound around an iron core. Small current magnetises the solenoid. This attracts to electrical contacts, making a complete circuit. Current flows from battery to starter motor.



Transverse wave	<i>Vibration causing the wave is at right angles to the direction of energy transfer</i>	Energy is carried outwards by the wave.	Water and light waves, S waves.
Longitudinal wave	<i>Vibration causing the wave is parallel to the direction of energy transfer</i>	Energy is carried along the wave.	Sound waves, P waves.

Electromagnet

Lots of turns of wire increase the magnetising effect when current flows

Turn current off, magnetism lost.

Increase strength of magnetic field

- Use larger current
- Use more turns of wire
- Put turns of wire closer together
- Use iron core in middle

Motor effect

Transverse and Longitudinal waves

Waves in air, fluids and solids

Properties

Wavelength	<i>Distance from one point on a wave to the same point of the next wave</i>
Amplitude	<i>The maximum disturbance from its rest position</i>
Frequency	<i>Number of waves per second</i>
Period	<i>Time taken to produce 1 complete wave</i>

Solenoid

A long coil of wire

Magnetic field from each loop adds to the next.

Right hand rule

Thumb: Direction of current.
Fingers: Direction of magnetic field.

Reverse current, magnetic field direction reverses.

Further away from the wire, magnetic field is weaker.

Current large enough, iron filings show circular magnetic field.

If current is small, magnetic field is very weak.

Electric current flowing in a wire produces a magnetic field around it.

AQA MAGNETISM **AQA Waves**

Electromagnetic waves

Electromagnetic wave *Continuous spectrum of transverse waves*

Wave speed	Wave speed = frequency X wavelength	$V = f \times \lambda$
Wave period	Wave period = $1 \div$ frequency	$T = 1 \div f$
Speed	Speed = distance \div time	$v = d \div t$

Permanent and Induced Magnetism

Magnets

Magnetic	<i>Materials attracted by magnets</i>	Uses non-contact force to attract magnetic materials.
North seeking pole	<i>End of magnet pointing north</i>	Compass needle is a bar magnet and points north.
South seeking pole	<i>End of magnet pointing south</i>	Like poles (N – N) repel, unlike poles (N – S) attract.
Magnetic field	<i>Region of force around magnet</i>	Strong field, force big. Weak field, force small. Field is strongest at the poles.
Permanent	<i>A magnet that produces its own magnetic field</i>	Will repel or attract other magnets and magnetic materials.
Induced	<i>A temporary magnet</i>	Becomes magnet when placed in a magnetic field.

Black surfaces	<i>Good emitters, good absorbers</i>
White surfaces	<i>Poor emitters, poor absorbers</i>
Shiny surfaces	<i>Good reflectors</i>



Properties

e.g. Gamma

Short wavelengths have high frequency and high energy.

	Units
Distance	<i>Metres (m)</i>
Wave speed	<i>Metres per second (m/s)</i>
Wavelength	<i>Metres (m)</i>
Frequency	<i>Hertz (Hz)</i>
Period	<i>Seconds (s)</i>

Specular	Flat surface reflection.
Diffuse	Rough surface reflection.

EM wave	Danger	Use
Radio	Safe.	Communications, TV, radio.
Microwave	Burning if concentrated.	Mobile phones, cooking, satellites.
Infrared		Heating, remote controls, cooking.
Visible	Damage to eyes.	Illumination, photography, fibre optics.
Ultra violet	Sunburn, cancer.	Security marking, disinfecting water.
X-ray	Cell destruction, mutation, cancer.	Broken bones, airport security.
Gamma		Sterilising, detecting and killing cancer.

